

Controller General of Patents, Designs & Trade
Marks



सत्यमेव जयते

G.A.R.6
[See Rule 22(1)]
RECEIPT



Docket No 10172

Date/Time 2024/01/29 19:05:06

To
Mahesh Madanrao Jadhav

UserId: maheshipr

Sr No 26/2/2/26, Yashodanand, Near Ram
Mandir, Ambegaon Pathar, Ambegaon (Bk),
Pune-411046

CBR Detail:

Sr. No.	App. Number	Ref. No./Application No.	Amount Paid	C.B.R. No.	Form Name	Remarks
1	202421005803	TEMP/E-1/6619/2024-MUM	1600	4198	FORM 1	"HERBAL FORMULATION FOR MANAGEMENT OF OSTEOARTHRITIS"

TransactionID	Payment Mode	Challan Identification Number	Amount Paid	Head of A/C No
N-0001313883	Online Bank Transfer	2901240044285	1600.00	1475001020000001

Total Amount : ₹ 1600.00

Amount in Words: Rupees One Thousand Six Hundred Only

Received from Mahesh Madanrao Jadhav the sum of ₹ 1600.00 on account of Payment of fee for above mentioned Application/Forms.

* This is a computer generated receipt, hence no signature required.

[Print](#)

[Home](#)

[About Us](#)

[Contact Us](#)



OPEN ACCESS

EDITED BY

Venkatakrishna Rao Jala,
University of Louisville, United States

REVIEWED BY

Robert Fultz,
Brightseed, United States
Alberto Finamore,
Council for Agricultural and Economics
Research (CREA), Italy
Dhrubajyoti Nag,
City College of New York (CUNY),
United States

*CORRESPONDENCE

Suresh Jagtap
✉ chiritatml@rediffmail.com

RECEIVED 04 August 2023

ACCEPTED 22 November 2023

PUBLISHED 08 December 2023

CITATION

Jadhav A, Jagtap S, Vyavahare S,
Sharbidre A and Kunchiraman B (2023)
Reviewing the potential of probiotics,
prebiotics and synbiotics: advancements
in treatment of ulcerative colitis.
Front. Cell. Infect. Microbiol. 13:1268041.
doi: 10.3389/fcimb.2023.1268041

COPYRIGHT

© 2023 Jadhav, Jagtap, Vyavahare, Sharbidre
and Kunchiraman. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Reviewing the potential of probiotics, prebiotics and synbiotics: advancements in treatment of ulcerative colitis

Apurva Jadhav¹, Suresh Jagtap^{1*}, Suresh Vyavahare²,
Archana Sharbidre³ and Bipinraj Kunchiraman⁴

¹Herbal Medicine, Interactive Research School for Health Affairs, Bharati Vidyapeeth (Deemed to be University), Pune, Maharashtra, India, ²Sai Ayurved Medical College, Maharashtra University of Health Sciences, Solapur, Maharashtra, India, ³Department of Zoology, Savitribai Phule Pune University, Pune, Maharashtra, India, ⁴Microbial Biotechnology, Rajiv Gandhi Institute of IT & Biotechnology, Bharati Vidyapeeth (Deemed to be University), Pune, Maharashtra, India

Inflammatory bowel diseases (IBD) like Crohn's and ulcerative colitis (UC) are multifactorial pathologies caused by environmental factors and genetic background. UC is a chronic inflammatory disorder that specifically targets the colon, resulting in inflammation. Various chemical interventions, including aminosalicylates, corticosteroids, immunomodulators, and biological therapies, have been extensively employed for the purpose of managing symptoms associated with UC. Nevertheless, it is important to note that these therapeutic interventions may give rise to undesirable consequences, including, but not limited to, the potential for weight gain, fluid retention, and heightened vulnerability to infections. Emerging therapeutic approaches for UC are costly due to their chronic nature. Alternatives like synbiotic therapy, combining prebiotics and probiotics, have gained attention for mitigating dysbiosis in UC patients. Prebiotics promote beneficial bacteria proliferation, while probiotics establish a balanced gut microbiota and regulate immune system functionality. The utilisation of synbiotics has been shown to improve the inflammatory response and promote the resolution of symptoms in individuals with UC through the stimulation of beneficial bacteria growth and the enhancement of intestinal barrier integrity. Hence, this review article aims to explore the potential benefits and underlying reasons for incorporating alternative approaches in the management of UC with studies performed using prebiotics, probiotics, and synbiotics to treat ulcerative colitis and to highlight safety and considerations in UC and future perspectives. This will facilitate the utilisation of novel treatment strategies for the safer and more efficacious management of patients with UC.

KEYWORDS

inflammatory bowel diseases, ulcerative colitis, probiotics, prebiotics, synbiotics, therapeutic uses

1 Introduction

Over the past decade, there has been an observed increase in the prevalence and incidence of Inflammatory Bowel Diseases (IBD), specifically Crohn's Disease (CD) and Ulcerative Colitis (UC) (Molodecky et al., 2012; Roy and Dhaneshwar, 2023). According to a recent epidemiological study conducted by Ng et al. (2017), IBD affects over 0.3% of people in North America and Europe. It is a group of gastrointestinal disorders with chronic inflammation. The cause of IBD is not fully understood but is believed to be a combination of genetic and environmental factors. Factors such as diet, use of antibiotics, and socioeconomic status can influence persistent immune-mediated inflammation in the intestines. (Manichanh et al., 2012; Çavdar and Çavdar, 2023). UC is a chronic gastrointestinal disorder characterised by chronic inflammation of the colorectal mucosa, resulting in non-specific inflammatory changes. The aetiology of this condition remains uncertain, and its conversion into cancer is not easily treatable. Ungaro et al. (2017) conducted epidemiological research which substantiated the global increase in the prevalence of UC. It is a condition caused by a combination of genetic, immune, microbial, and environmental factors. It involves persistent inflammation of the mucosal lining, affecting the rectum and proximal colon segments. The exact mechanisms behind UC development are not fully understood, but it is widely recognised that environmental factors, host genetics, a dysregulated immune system, and microbial dysbiosis play crucial roles. (Ananthkrishnan, 2015).

According to the study conducted by Higuchi et al. (2012), there is evidence to suggest that the adoption of Westernised diets and lifestyles can heighten the susceptibility of genetically predisposed individuals to IBD. Environmental factors encompass various aspects, such as residing in industrialised nations, urbanised settings, and regions situated at higher latitudes. In addition to infections and smoking, other factors also play a role in the pathogenesis of IBD. Patients frequently need lifelong medication, which is the main purpose of therapy, both to persuade clinical remission and to conserve it for a long period. According to the Global Burden of Disease Study, the estimated global prevalence of Crohn's disease was approximately 6.8 million individuals, while ulcerative colitis affected approximately 10 million individuals. (Alatab et al., 2020).

These data suggest that, in addition to finding effective therapeutic approaches to advance IBD patients' quality of life and lower healthcare system costs, it is crucial to better understand the factors involved in the development of IBD. Costs of IBD management have increased significantly in the past five years due to different therapeutic interventions and disease characteristics. Patients face increased healthcare utilisation, out-of-pocket expenses, and reduced productivity, necessitating the implementation of cost-effective solutions. (Park et al., 2020) Recent research suggests that microbiota can be therapeutically effective in treating IBD. The gut microbiome influences host processes and can mitigate inflammation. Various therapeutic approaches, including prebiotics, probiotics, and synbiotics, have been developed to alter and restructure the gut microbiome.

(Britton et al., 2021) The most common strains currently available as probiotics and possessing beneficial health effects are *Enterococcus faecium*, *Bifidobacterium*, *Bacillus*, *Saccharomyces boulardii* (*S. boulardii*) and *Lactobacillus* strains. Prebiotics like fructo-oligosaccharides (FOS), isomalto-oligosaccharides (IMO), and xylooligosaccharide (XOS) are studied for their potential benefits on stool volume, constipation relief, and faecal acidity, with easily metabolised by *Bifidobacteria* and other microorganisms. (Guarino et al., 2020).

2 Understanding ulcerative colitis

Ulcerative colitis is a chronic inflammatory disorder of the gastrointestinal tract characterised by a progressive decline in health. UC is characterised by inflammation of the mucosal lining, typically confined to the colon and rectum. On the other hand, CD involves inflammation that extends through the entire thickness of the gastrointestinal tract, with a predilection for the ileum, terminal ileum, and colon. The CD is often associated with additional inflammatory conditions outside of the intestines. According to Çavdar and Çavdar (2023), the medical characteristics encompass symptoms such as hemorrhagic diarrhoea, weight loss, fatigue, and abdominal pain. Additionally, certain individuals may develop extra-intestinal manifestations associated with IBD, such as primary sclerosing cholangitis, skin lesions, or joint complications (Judge and Lichtenstein, 2001). The impact of UC on a global scale has been evaluated, revealing that it affects a population of approximately 5 million individuals. UC is characterised by recurring episodes of inflammation in the mucosal layer of the colon, which alternate between periods of relapse and remission. In ulcerative colitis, inflammation is typically confined to the mucosal surface. The disorder originates in the rectum and typically progresses proximally throughout the entirety of the colon. To date, a conclusive therapeutic approach for UC remains elusive, and the currently available interventions primarily focus on sustaining remission (Roy and Dhaneshwar, 2023).

Conventional therapies for UC consist of pharmacological interventions involving the use of various agents, including azathioprine, aminosalicylates, and corticosteroids. These treatments aim to induce remission, prevent relapses, promote mucosal healing, and ultimately reduce the need for colectomy in individuals diagnosed with UC. In recent times, advancements in medical treatments have been influenced by the utilisation of agents such as monoclonal antibodies that specifically target proinflammatory cytokines, adhesion molecules, T-cell activation, and agents that promote the production of anti-inflammatory cytokines, such as IL-10 (Interleukin) and Transforming Growth Factor (TGF). However, the administration of these medications in the context of ulcerative colitis is associated with a distinct set of systemic and localised adverse effects experienced by individuals. These include but are not limited to, symptoms such as headache, abdominal pain, nausea, cramping, loss of appetite, vomiting, and rash. In addition, it is not recommended to use corticosteroids for an extended period due to their potential to induce glucose

intolerance, osteoporosis, myopathy, and increased vulnerability to infections. These adverse effects raise significant concerns regarding the safety profile of corticosteroid usage (Mowat et al., 2011). Similarly, the expenses associated with these promising therapeutic approaches are substantial, considering the increasing prevalence and persistent nature of IBD. Histological examinations conducted on patients with UC have revealed the presence of impaired intestinal epithelial barrier function. However, the significance of this dysfunction with the pathology of UC remains uncertain. Patients diagnosed with UC exhibit changes in the function of the epithelial barrier, impaired function of tight junctions, and reduced secretion of mucin-2. Nevertheless, the exact relationship between these abnormalities and chronic inflammation, accompanied by heightened cytokine production, remains uncertain. Heller et al. (2005) conducted a study that yielded evidence indicating that the overexpression of IL-13 plays a pivotal role in the disruption of tight junctions within the intestinal mucosal layer of individuals who have been diagnosed with active UC. The dysfunction of the barrier function in individuals diagnosed with UC has been linked to a heightened absorption of luminal antigens and an elevated susceptibility to bacterial infiltration in the underlying mucosa. Extensive research has provided comprehensive insights into the dysregulation of the intestinal immune system in individuals afflicted with UC. Dendritic cells (DCs) are known to have a pivotal role in the augmentation of UC patients, thus signifying the substantial contribution of these cells in the initiation and perpetuation of inflammatory processes (Hart et al., 2005).

3 Conventional treatments for UC

Conventional therapies play a pivotal role in the management of ulcerative colitis and the mitigation of associated symptoms. The objective of these treatments is to manage inflammation, initiate and sustain remission, and enhance the overall quality of life for patients. The Selecting Therapeutic Targets in IBD (STRIDE)-II statement was revised by the International Organisation for the Study of IBD to enhance the treatment of IBD in adults and children. The revised statement identified several goals at different timeframes. In the short term, achieving symptomatic remission and normalising C-reactive protein levels were prioritised. In the intermediate term, reducing calprotectin levels was identified as a target. Lastly, in the long term, the goals included achieving endoscopic healing and standardising the quality of life for individuals with IBD (Im et al., 2018; Turner et al., 2021). The subsequent therapies listed are commonly employed conventional treatments for ulcerative colitis.

3.1 Aminosalicylates

Aminosalicylates, alternatively referred to as 5-aminosalicylic acid (5-ASA) drugs, encompass a class of pharmaceuticals that possess the capability to mitigate inflammation within the gastrointestinal tract. These therapeutic agents function by

specifically targeting the mucosal lining of the colon and rectum, which is the site of inflammation. Aminosalicylates are offered in diverse formulations, encompassing oral pharmaceuticals, rectal suppositories, and enemas. Corticosteroids are frequently employed as an initial therapeutic approach for individuals with mild to moderate ulcerative colitis, effectively managing symptoms and sustaining periods of remission. The class of medications known as 5-ASA-based drugs includes sulfasalazine, olsalazine, Mesalazine and balsalazide. These medications are widely recognised for their efficacy, safety, and affordability in the treatment of IBD, particularly UC (Sood et al., 2019).

3.2 Corticosteroids

Corticosteroids, such as prednisone and budesonide, exhibit potent anti-inflammatory properties and have demonstrated efficacy in promptly alleviating symptoms associated with moderate to severe ulcerative colitis exacerbations. These medications function through the inhibition of the immune system and the mitigation of inflammation within the colon. Corticosteroids are commonly prescribed for a limited duration to induce remission or manage acute symptoms. The avoidance of prolonged corticosteroid use is typically recommended due to the potential for significant adverse effects. According to Wild et al. (2003) and Waljee et al. (2016), these interventions effectively decrease intestinal inflammation by promptly reducing intestinal permeability, suppressing the production of Tumour necrosis factor (TNF) and inhibiting Nuclear factor- κ B (NF- κ B).

3.3 Immunomodulators

Immunomodulators represent a category of pharmaceutical agents that exert a modulatory or regulatory effect on the immune system. Patients who exhibit poor responses to aminosalicylates or corticosteroids are frequently prescribed these medications. Additionally, they may be used as a maintenance therapy to mitigate the occurrence of flare-ups. mercaptopurine (MP), Azathioprine (AZT) and methotrexate (MTX) are among the immunomodulators commonly employed in the treatment of ulcerative colitis. These pharmaceutical agents function by inhibiting the immune response and diminishing inflammation within the colon. According to Singh et al. (2022), in cases where patients with IBD do not respond to 5-ASA drugs and are reliant on or unresponsive to corticosteroids, it is advised to utilise conventional immunomodulators such as 6-MP, AZT, and MTX to maintain remission.

3.4 Biologic therapies

Biologic therapies are a relatively new class of pharmaceutical interventions that specifically target specific molecules or cells involved in the inflammatory pathway of ulcerative colitis. These interventions are usually recommended for people with moderate-

to-severe symptoms who have not responded to conventional treatment. Biological therapeutics, such as anti-tumour necrosis factor (TNF) agents (e.g., adalimumab & infliximab) or integrin receptor antagonists (e.g., vedolizumab), function by inhibiting specific proteins or cells involved in the inflammatory processes taking place in the gastrointestinal tract. In contemporary medical practice, infliximab and adalimumab are the prevailing anti-tumour necrosis factor-alpha (TNF- α) agents that are frequently employed. The administration of these medications occurs through intravenous and subcutaneous routes, respectively. According to Seo and Chae (2014), clinical trials have provided evidence supporting the efficacy of infliximab and adalimumab in the treatment of moderate-to-severe UC.

3.5 Surgical interventions

Surgical interventions are necessary for treating ulcerative colitis when severe symptoms arise, complications arise, or conventional treatments are ineffective. Proctocolectomy involves removing the colon and rectum, with or without an ostomy. This procedure has high efficacy in eradicating the afflicted colon, providing patients with a prolonged period of remission. Treatment selection depends on the severity and scope of ulcerative colitis, as well as individual patient factors and preferences. (Kuhn and Klar, 2015).

4 Antibiotics used in UC treatment

Rifaximin modifies gut microbiota for small intestinal bacterial overgrowth and hepatic encephalopathy, while antibiotics aim to manipulate intestinal flora, potentially disrupting IBD progression. In their study, Khan et al. (2011) conducted a meta-analysis that examined the findings of 10 randomised controlled trials (RCTs) focused on the utilisation of antibiotics in patients diagnosed with active CD. Additionally, the study also included nine RCTs that investigated the impact of antibiotic usage in individuals with UC. Antibiotics significantly impact remission and disease severity in UC and CD. Rifamycin derivatives are effective in treating active CD. However, results vary due to different trials involving rifaximin, fluoroquinolones, macrolides, and anti-tuberculosis antibiotics. (Steinhart et al., 2002).

5 The role of gut microbiota in UC

The gut microbiota, an important group of microorganisms that live in the digestive tract, is correlated to several diseases, including ulcerative colitis. The composition and function of the microbiota in the gut have both been shown to be susceptible to change, as recent research has shown. The gastrointestinal tract of a neonate does not initially contain any microorganisms but is eventually colonised by bacteria that come from the mother and the environment. (Dominguez-Bello et al., 2010). According to Vyas and Ranganathan (2012), the number of bacterial cells present on

and within the entire body of an adult human is approximately ten times greater than the total number of human cells. The complexity and diversity of the human microbiome are considerable. The composition and quantity of anatomical features vary across different regions of the gastrointestinal tract, ranging from the nasal and oral cavities to the distal segments of the colon and rectum. The composition and complexity of the gut microbiota exhibit variability during the transition of infants from a liquid diet to solid foods. The alterations in dietary patterns during adulthood play a substantial role in shaping the composition of the gut microbiota. Metagenomic techniques based on 16S ribosomal RNA sequence have improved our understanding of microbial communities in the gastrointestinal tract. This approach reveals that 90% of the bacterial population is attributed to Bacteroidetes and Firmicutes. (Mariat et al., 2009; Hu and Gubatan, 2023).

5.1 Dysbiosis in UC

Dysbiosis in the gut microbiota is the root cause of ulcerative colitis. This condition is characterised by a decrease in commensal bacteria such as bifidobacteria and *Lactobacilli* and an increase in pathogenic bacteria such as *Escherichia coli* and *Fusobacterium*. The imbalance that can result from dysbiosis can put a person at risk for several different diseases, including IBD and irritable bowel syndrome. Additionally, it has the potential to bring about unanticipated outcomes, such as the activation of an HIV infection or the development of autoimmune diseases. (Liu et al., 2016).

5.2 Impaired gut barrier function

The gut microbiota and intestinal epithelial cells collaborate to establish a defensive barrier aimed at impeding the infiltration of pathogens and toxins into the bloodstream. The integrity of the barrier function in ulcerative colitis is compromised, resulting in heightened permeability of the intestinal lining. The presence of dysbiosis in individuals with ulcerative colitis has been observed to have a detrimental effect on the integrity of the gut barrier. This compromised barrier allows for the infiltration of pathogenic bacteria and their associated metabolic by-products into the intestinal mucosa, subsequently eliciting an immune response. This phenomenon gives rise to atypical immune-inflammatory responses, including inflammation, allergies, and autoimmune disorders, which are facilitated by molecular mimicry and dysregulated T-cell reactions. According to the study conducted by Barnaba and Sinigaglia, 1997, it was found that the maintenance of intestinal homeostasis and the differentiation between harmful pathogens and beneficial commensal microbes heavily rely on the Treg/TH17 ratio, which represents the equilibrium between regulatory T cells (Tregs) and T helper type 17 cells (TH17). This regulation is strongly linked to the gut microbiota, as commensal microorganisms, including *Firmicutes*, *Bacteroides fragilis* and *Bifidobacterium infantis* promote the expansion of Treg cells, specifically FOXP3-expressing Treg and IL-10-producing Treg

lymphocytes. Foxp3+ regulatory T cells play a crucial role in gut immunity and physiology, inducing intestinal tolerance and defending the host against harmful dietary antigens, commensal microorganisms, and pathogens. They also facilitate local tissue repair and maintain the integrity of the epithelial barrier, making them essential non-immune cells in the gastrointestinal tract. (Cosovanu and Neumann, 2020).

5.3 Immune system dysregulation

The gut microbiota exerts a substantial impact on the maturation and modulation of the immune system. Individuals diagnosed with UC exhibit an atypical immune response towards the gut microbiota, which is distinguished by an intensified activation of the immune system and the persistence of inflammatory processes. The dysregulation of the immune system can sustain the ongoing cycle of inflammation and subsequent tissue damage in the colon, thereby playing a role in the advancement of ulcerative colitis. The dysregulation of the immune response in the intestinal region is a crucial factor in the development of IBD, involving various molecules such as cytokines. In a recent study conducted by Kmiec et al. (2017), it was demonstrated that there is a correlation between the innate immune response and the promotion of gut inflammation in patients with IBD. The mucosa of patients with UC exhibits an altered equilibrium between regulatory T-cells and effector T-cells, including Th1, Th2, and Th17 cells. Evidence suggests that there is a correlation between UC and an atypical type 2 immune response, which is mediated by non-classical natural killer T-cells that produce IL-5 and IL-13. Previous studies have demonstrated that IL-13, which is secreted by specific subsets of NKT cells, plays a crucial role in exerting cytotoxic effects on epithelial cells. These effects include the induction of apoptosis and alterations in the composition of tight junction proteins (Fuss and Strober, 2008; Heller et al., 2008). Poggi et al. (2019) demonstrated a correlation between the exacerbation of pathology and the presence of supplementary inflammatory cytokines, including TNF, IL-1, IL-6, and IL-9.

5.4 Role of short-chain fatty acids

Short-chain fatty acids (SCFAs) are one of many metabolites produced by the gut microbiota during the fermentation of dietary fibres, exhibit anti-inflammatory properties. These fatty acids, typically not exceeding six carbon atoms, are derived from prebiotic or microbially fermentable carbohydrates like inulin, polysaccharides, and resistant starch. (Dalile et al., 2019) Butyrate, acetate, and propionate are examples of SCFAs that play important roles in intestinal health. The observed effects include acceleration of the regeneration and healing process of intestinal epithelial cells, augmentation of mucus production, and maintenance of appropriate pH levels within the intestine. Moreover, they hinder the attachment of pathogenic microorganisms to enterocytes. Acetate is commonly utilised as a cellular energy substrate for the

growth and maintenance of muscle tissue and colonic cells. Butyrate exhibits a variety of beneficial effects on the host, such as strengthening metabolism, modulating the host's immune system, and facilitating anti-inflammatory mechanisms. As a result, it attracts considerable attention. (Guarino et al., 2020) One of the hallmarks of UC is a decrease in SCFAs production, which contributes to an already severe inflammatory response and jeopardises the GI tract's overall health. Because of this decline in SCFAs producers, it is common to find lower levels of SCFAs in faeces taken from people with inflammatory IBD. In a study of people with UC, Machiels et al. (2013) found lower levels of acetate and propionate in their faeces but not butyrate. When studying faecal samples from people with IBD Huda-Faujan et al. (2010) found lower levels of butyrate and propionate. Recent research findings indicate notable disparities in gut microbial species, microbial diversity and metabolic pathways between individuals diagnosed with IBD and healthy individual. (Lavelle and Sokol, 2020)

5.5 Potential therapeutic strategies

Due to the significant impact of the gut microbiota on ulcerative colitis, there is a growing interest in the development of effective strategies aimed at modulating the microbiota to restore equilibrium and improve disease outcomes. This involves the utilisation of probiotics, which are live beneficial bacteria that can be employed to reinstate the equilibrium of microorganisms in the gastrointestinal tract. Prebiotics, classified as dietary fibres that possess the ability to selectively enhance the proliferation of advantageous bacteria, can additionally be employed to enhance the growth of beneficial microorganisms. In addition, the utilisation of synbiotics, a combination of probiotics and prebiotics, may exhibit synergistic outcomes in the restoration of equilibrium within the gut microbiota (Hu and Gubatan, 2023). The present review provides an overview of the existing microbiome-centred therapeutic strategies employed in the management of IBD, as outlined in Table 1.

6 Probiotics: restoring balance in the gut

Probiotics can be traced back to ancient civilisations when humans first began eating fermented foods as part of their regular diet. It was hypothesised by Elie Metchnikoff that the beneficial bacteria found in yoghurt could improve the health of the gut microbiome. Mackowiak (2013) presented the concept of probiotics for the first time. These microorganisms can positively modulate the composition of the gut microflora and reduce the presence of pathogenic bacteria that release harmful compounds within the human gastrointestinal tract. In previous studies, a range of microorganisms have been employed for disease management, thereby giving rise to the term "probiotics". Probiotics, derived from Greek, refer to living, non-pathogenic organisms with


TABLE 1 Summary of current microbiome-based therapeutic strategies in IBD.

Therapeutic	Examples	Effects on microbiome
Antibiotics	Rifaximin Ciprofloxacin Metronidazole Tobramycin Amoxicillin	<ul style="list-style-type: none"> > Eliminate specific microbial populations that contribute to hyperinflammatory states to modulate IBD severity/disease activity > Prevent overgrowth of harmful microbial species that may lead to secondary IBD complications (e.g. pouchitis and abscesses) > Influence the development of anti-drug antibodies that affect the risk of immunogenicity to anti-TNF biologics > Increase risk of microbial resistance > Increase risk of infections (e.g. <i>Clostridium difficile</i>)
Prebiotics (molecular compounds)	Lactulose Psyllium Fructo-oligosaccharides Germinated barley foodstuff	Metabolised by gut microorganisms to form small molecule by-products (e.g. butyrate and short-chain fatty acids) that influence the local microenvironment to preferentially favour growth of certain flora
Probiotics (living microorganisms)	<i>Lactobacillus</i> spp. <i>Bifidobacterium</i> spp. <i>Saccharomyces</i> spp. Non-pathogenic <i>Escherichia coli</i>	<ul style="list-style-type: none"> > Strengthen intestinal barrier function by inhibiting apoptosis of intestinal cells > Regulate immunity through genetic pathways (NF-κB, IL-6, TNF-α) or direct influence on T-cells > Produce small molecules (lactic acid, hydroperoxides) that directly influence the growth patterns of other microbial strains Provide local survival competition for scarce resources with other microorganisms
Synbiotics	Combination of prebiotics and probiotics	<ul style="list-style-type: none"> > Optimise a combination of prebiotics and probiotics for maximum synergistic effect > Prebiotic(s) are specifically chosen to select for the growth and survival of probiotic organism(s)

beneficial effects on hosts. Vergin, in 1954, coined the term “probiotic” to compare the harmful effects of antibiotics on intestinal microbiota with the beneficial effects of some beneficial bacteria. The term “probiotic” was subsequently redefined by Lilly and Stillwell as a biologically active product produced by a microorganism that enhances the growth of another microorganism. As a result, the term was subsequently defined by Fuller (1989) as non-pathogenic microorganisms that, when ingested, have a beneficial impact on the health or physiology of the host. The most recent definition proposed by the FDA and WHO in collaboration is that of a live microorganism that, when administered in adequate quantities, confers a health advantage to the host. Several probiotic microorganisms are commonly utilised, species of *Bifidobacterium* and *Lactobacillus*, have been reported to be the predominant and subdominant groups among gut microbes, respectively (Casellas et al., 2007; Markowiak and Śliżewska, 2017). Bacterial organisms capable of forming spores, primarily belonging to the genus *Bacillus*, are prevalent in this context (Derikx et al., 2016; Jin et al., 2023).

Probiotics are incorporated into various food products, particularly fermented dairy products, either individually or in conjunction with other strains. Continuously, novel genera and strains of probiotics are being discovered through increasingly sophisticated and targeted research endeavours. VSL#3 is a combination of eight distinct probiotic strains, exhibiting high strain specificity and variability in benefits across different patient groups. Probiotic products can consist of a solitary strain or a mixture of two or more strains, with the effectiveness of multi-strain probiotics demonstrated in controlled studies. (Chapman et al., 2011). The study of probiotics, particularly *Lactobacilli*, has experienced significant growth over the past twenty years. This is evident from the substantial increase in research articles published on the topic. Specifically, while there were approximately 180 research articles available between 1980 and 2000, the number of articles on the probiotic *Lactobacillus* exceeded 5700 between 2000 and 2014. According to a study published in PubMed in 2014, the effectiveness of the probiotic strain *Lactobacillus* was investigated. The characteristics of an ideal probiotic strain are depicted in Figure 1.

Probiotics act against intestinal diseases through various processes, including colonisation, which enhances their efficacy. They produce inhibitory constituents like organic acids, fatty acids, hydrogen peroxide, SCFAs, and bacteriocin-like inhibitory substances that prevent pathogens. (Tharmaraj and Shah, 2009). Bacteriocins refer to antimicrobial peptides or proteins that are produced by bacteria via ribosomal mechanisms. The aforementioned compounds demonstrate a diverse spectrum of antimicrobial efficacy against pathogenic bacteria, encompassing multiple strains of *Staphylococcus*, *Listeria*, *Bacillus*, *Clostridium*, and other bacterial species (Dai et al., 2021; Fathizadeh et al., 2022). The significance of bacteriocins synthesised by probiotic bacteria is ascribed to their secure utilisation in diverse domains, including the food industry, pharmaceuticals, veterinary medicine, and human healthcare (Barcenilla et al., 2021; Sharma et al., 2021; Somashekaraiah et al., 2021). The compounds cited in the scholarly articles (Sheoran and Tiwari, 2019; Sheoran and Tiwari, 2020; Chen et al., 2021) exhibit inhibitory effects on a wide range of bacteria. Their mechanism of action involves reducing cell viability, modifying bacterial cell metabolism, and inhibiting toxin production. Additional mechanisms of probiotic action include competitive inhibition on the surface of the intestinal epithelium, wherein the probiotics obstruct the binding sites of the intestinal epithelial surface, as well as a reduction in the interaction between the pathogen and the host. Similarly, the investigation of nutrient competition is also examined as a mechanism underlying probiotic activity. According to the suggestions of the World Health Organization (WHO), Food and Agriculture Organization (FAO), and the European Food Safety Authority (EFSA), in their selection process, probiotic strains must meet both safety and functionality criteria, as well as those related to their technological usefulness. Probiotic characteristics are not associated with the genus or species of a microorganism, but with few and specially selected strains of a particular species (Hill et al., 2014). FAO and WHO have jointly developed guidelines that propose a systematic approach for conducting comprehensive evaluations of probiotics in food



products, to provide scientific evidence to support health claims and benefits. The essential criteria for an ideal probiotic are presented by Pandey et al. (2015). The application of the FAO/WHO guidelines on Probiotics has the potential to serve as a universal benchmark for evaluating probiotics in food products, thereby facilitating the validation of health-related assertions.

The guidelines stipulate that the following activities must be carried out:

The identification and categorisation of strains.

- A comprehensive elucidation of the strain(s) about their safety and probiotic attributes.
- Verification of the health benefits observed in human studies.
- Ensuring the accuracy and transparency of efficacy claims and product information throughout the entire duration of the product's shelf life.

6.1 Probiotics in ulcerative colitis

The colon contains the highest concentration of microorganisms, suggesting treating colon microbiome abnormalities could benefit individuals with ulcerative colitis. Studies have shown potential benefits from various probiotic strains. (Basso et al., 2019). Previous studies have demonstrated that the use of the non-pathogenic strain of *E. coli* Nissle 1917 (EcN) exhibited similar efficacy and safety characteristics in the maintenance therapy of patients with mild or moderate ulcerative colitis when compared to treatment with salicylates (Basso et al., 2019; Bischoff et al., 2020). The potential therapeutic and prophylactic effects of the probiotic strain EcN against adherent-invasive *Escherichia coli* (AIEC) infection in zebrafish were investigated. In the zebrafish model, the researchers discovered that EcN effectively reduced AIEC colonisation, tissue damage, and pro-inflammatory responses. Furthermore, EcN reduced AIEC hyperinfection in zebrafish, especially when

propionic acid was present. The effectiveness of EcN in combating AIEC infection in a zebrafish model is highlighted in this study. (Nag et al., 2022) The utilisation of various strains of lactic acid bacteria and bifidobacteria as supplementary treatment has been observed to significantly improve the progression of the disease and the sustained absence of symptoms in individuals diagnosed with ulcerative colitis (Basso et al., 2019). A highly regarded probiotic complex is VSL#3, which consists of four strains of *Lactobacillus* (*L. casei*, *L. acidophilus*, *L. plantarum* and *L. delbrueckii* subsp. *bulgaricus*), three strains of *Bifidobacterium* (*B. longum*, *B. breve*, and *B. infantis*), and one strain of *Streptococcus* (*S. salivarius* subsp. *termophilus*) (Basso et al., 2019). Studies conducted on mouse models have demonstrated that the administration of this probiotic mixture leads to the suppression of NF- κ B and TNF expression in the TLR4-NF- κ B signalling pathway. Consequently, the downregulation of pro-inflammatory cytokines and toll-like receptors (TLR) occurs while the upregulation of regulatory cytokines is observed (Jakubczyk et al., 2020; Silva et al., 2020). Previous studies have demonstrated the efficacy of VSL#3 in inducing and sustaining remission in individuals with mild to moderate UC when used as adjuvant therapy or as a standalone treatment (Basso et al., 2019).

In a systematic review and meta-analysis conducted by Derwa et al. (2017), it was found that the VSL#3 probiotic mixture demonstrated potential efficacy in inducing remission among individuals with UC. Furthermore, the study suggested that the use of VSL#3 probiotics may be comparable to 5-ASA in terms of preventing disease exacerbations. Under the recommendations put forth by the European Society for Clinical Nutrition and Metabolism (ESPEN), the investigation of probiotics such as VSL#3 and EcN in the treatment of mild to moderate UC is warranted, as they have shown potential for inducing remission in affected patients. According to Bischoff et al. (2020), the use of probiotics is contraindicated in cases of severe ulcerative UC. The use of probiotics as a supplementary treatment may have particular efficacy in the management of patients who experience intolerance to 5-ASA (Miele et al., 2018; Bischoff et al., 2020). Table 2 presents

several clinical trial findings regarding the use of probiotics for the treatment of UC.

7 Prebiotics: nourishing beneficial bacteria

The term “prebiotics” is a relatively recent concept that is clearly defined as “nondigestible food ingredients that beneficially alter the host by selectively stimulating the growth and/or activity of a specific group of bacteria in the colon, thereby improving the overall health of the host” (Gibson and Roberfroid, 1995). Recently, the International Scientific Association for Probiotics and Prebiotics (ISAPP) convened a panel of experts to reassess the definition of prebiotics. The panel has expanded the definition to encompass a broader understanding, defining it as “a substrate that is selectively utilised by host microorganisms, resulting in a health benefit.” The definition presented by Gibson et al. (2017) expands upon the concept of prebiotics by considering the potential inclusion of non-carbohydrate substances, exploring their relevance to body sites beyond the gastrointestinal tract, and identifying distinct groups beyond those found in food. These substances are typically classified as short-chain carbohydrates that are resistant to digestion yet serve as substrates for the proliferation of probiotic microorganisms within the upper gastrointestinal tract. Cocoa-derived flavanols are compounds that are not classified as carbohydrates but have been proposed as potential prebiotics. According to Tzounis et al. (2011), experimental studies conducted both *in vivo* and *in vitro* demonstrate that flavanols can enhance the proliferation of lactic acid bacteria.

A significant increase in the *Bifidobacteria* population in faecal samples has been reported due to fructooligosaccharides (FOS) consumption. Prebiotics, consisting of glucose, fructose, galactose and/or xylose, undergo minimal hydrolysis in the intestinal tract and have caloric value due to their resistance to digestion and energy metabolism through fermentation (Roberfroid, 1993). Prebiotics like FOS, isomalto-oligosaccharides (IMO), and xylooligosaccharides (XOS) are studied for their potential benefits on stool volume, constipation relief, and faecal acidity. FOS, like inulin and neosugar, are dietary fibres that improve stool volume and faecal acidity. These prebiotics are easily metabolised by *bifidobacteria* and other microorganisms like *Lactobacillus acidophilus*, *Bacteroides vulgatus*, *B. ovatus*, *B. thetaiotaomicron*, *B. fragilis*, and *Enterococcus faecium*, *E. faecalis* (Guarino et al., 2020) An optimal prebiotic substance should possess several key characteristics, as illustrated in Figure 2. IMO can be identified in various fermented food products, including miso, soy sauce, and honey. These compounds are known to be metabolised by *bifidobacteria* and the *Bacteroides* groups. IMO facilitates the proliferation of *Bifidobacterium* and *Lactobacillus* species, leading to both local and systemic Th1-like immune responses and the regulation of immune function. The presence of positive outcomes has been identified in the clinical trials. XOS are a class of prebiotics that occur naturally in various sources such as fruits, bamboo shoots, vegetables, milk, honey, and others (Aachary and Prapulla, 2010). Previous research

TABLE 2 Some clinical trial data of probiotics for treating UC.

Sr. no.	Probiotic Used	Outcome	Reference
1	A probiotic product that contained <i>L. casei</i> Zhang, <i>L. plantarum</i> P-8 and <i>B. animalis</i> subsp. <i>lactis</i>	The overall remission rate was 91.67% for the probiotic group vs. 69.23% for the placebo group ($P = 0.034$)	Chen et al., 2020a.
2	Symprove (contains <i>Lactobacillus rhamnosus</i> NCIMB 30174, <i>Lactobacillus plantarum</i> NCIMB 30173, <i>Lactobacillus acidophilus</i> NCIMB 30175 and <i>Enterococcus faecium</i> NCIMB 30176)	The calprotectin levels were significantly decreased following 4 weeks in the probiotic group ($p = 0.011$ and 0.001 , t-test and Wilcoxon's, respectively)	Bjarnason et al., 2019.
3	Kefir (<i>Lactobacillus</i> Bacteria)	No statistically significant difference was found between weeks 1 and 2 in patients with UC in terms of abdominal pain, bloating, frequency of stools, defecation consistency, and feeling good.	Yilmaz et al., 2019.
4	A tablet contains <i>Streptococcus faecalis</i> T-110, <i>Clostridium butyricum</i> TO-A and <i>Bacillus mesentericus</i> TO-A	At 12 months, the remission rate was 69.5% in the treatment group and 56.6% in the placebo group ($p = 0.248$). The relapse rates in the treatment and placebo groups were 0.0% vs. 17.4% at months ($p = 0.036$).	Yoshimatsu et al., 2015.
5	Mil-Mil (a fermented milk product containing <i>B. breve</i> strain Yakult and <i>Lactobacillus acidophilus</i>)	Relapse-free survival was not significantly different between the treatment and placebo groups ($P = 0.643$)	Matsuoka et al., 2018.
6	<i>Escherichia coli</i> Nissle 1917	Efficacy in maintaining remission and preventing relapse comparable to Mesalazine	Kruis et al., 2004.
7	<i>Escherichia coli</i> Nissle 1917	Efficacy in maintaining remission after exacerbation of UC comparable to mesalazine	Rembacken et al., 1999.
8	<i>Escherichia coli</i> Nissle 1917	Efficacy and safety in maintaining remission comparable to mesalazine	Kruis et al., 2004.
9	<i>Escherichia coli</i> Nissle 1917	Efficacy in maintaining remission comparable to mesalazine	Henker et al., 2008.
10	<i>Escherichia coli</i> Nissle 1917	Possibility of dose-dependent efficacy in inducing remission of	Matthes et al., 2010.

(Continued)

TABLE 2 Continued

Sr. no.	Probiotic Used	Outcome	Reference
		the rectal probiotic compared to placebo	
11	<i>Escherichia coli</i> Nissle 1917	No benefit in the use of probiotics as an additional therapy to conventional treatment	Petersen et al., 2014.
12	<i>Lactobacillus</i> GG	Higher efficacy of probiotics as add-on therapy in prolonging the relapse-free time compared to mesalazine monotherapy	Zocco et al., 2006.
13	<i>Bifidobacterium breve</i> , <i>Bifidobacterium bifidum</i> , <i>Lactobacillus acidophilus</i> YIT 0168 (<i>Bifidobacteria</i> -Fermented Milk- BFM)	Higher efficacy of probiotic mixture as add-on therapy in maintaining remission and preventing relapse compared to conventional therapy alone	Ishikawa et al., 2003.
14	<i>Bifidobacterium breve</i> , <i>Bifidobacterium bifidum</i> , <i>Lactobacillus acidophilus</i> YIT 0168 (<i>Bifidobacteria</i> -Fermented Milk- BFM)	Higher efficacy of probiotics as add-on therapy in maintaining remission compared to conventional therapy alone	Kato et al., 2004.
16	<i>Lactobacillus reuteri</i> ATCC 55730	Higher efficacy of probiotic enema as add-on therapy additional to oral mesalazine in improving mucosal inflammation compared to conventional therapy	Oliva et al., 2011.
17	<i>Lactobacillus casei</i> , <i>Lactobacillus plantarum</i> , <i>Lactobacillus acidophilus</i> and <i>Lactobacillus delbrueckii</i> subsp. <i>Bulgaricus</i> , <i>Bifidobacterium longum</i> , <i>Bifidobacterium breve</i> and <i>Bifidobacterium infantis</i> , <i>Streptococcus salivarius</i> subsp. <i>Thermophilus</i> (VSL#3)	The higher efficacy of probiotic mixture as add-on therapy to conventional treatment in patients with the relapsing disease compared to placebo	Tursi et al., 2010.
18	<i>Lactobacillus casei</i> , <i>Lactobacillus plantarum</i> , <i>Lactobacillus acidophilus</i> and <i>Lactobacillus delbrueckii</i> subsp. <i>Bulgaricus</i> , <i>Bifidobacterium longum</i> , <i>Bifidobacterium breve</i> and <i>Bifidobacterium infantis</i> , <i>Streptococcus salivarius</i> subsp. <i>Thermophilus</i> (VSL#3)	Higher efficacy in inducing and maintaining remission compared to placebo	Sood et al., 2009.

(Continued)

TABLE 2 Continued

Sr. no.	Probiotic Used	Outcome	Reference
19	<i>Lactobacillus casei</i> , <i>Lactobacillus plantarum</i> , <i>Lactobacillus acidophilus</i> and <i>Lactobacillus delbrueckii</i> subsp. <i>Bulgaricus</i> , <i>Bifidobacterium longum</i> , <i>Bifidobacterium breve</i> and <i>Bifidobacterium infantis</i> , <i>Streptococcus salivarius</i> subsp. <i>Thermophilus</i> (VSL#3)	Higher efficacy in maintaining remission compared to placebo	Miele et al., 2009.
20	<i>Lactobacillus plantarum</i> , <i>Bifidobacterium longum</i> , and <i>Bifidobacterium bifidum</i>	Suppressed colonic inflammation, and fatigue by the suppression of the IL-1 β or IL-6 to IL-10 expression ratio and gut bacterial LPS production	Yoo et al., 2022.
21	<i>Lactobacillus plantarum</i>	Restored gut microbiota balance and modulated the resident gut microbiota and immune response	Khan et al., 2022.
22	<i>Ligilactobacillus salivarius</i> Li01 and RSV	An improved synergistic anti-inflammatory effect from the RSV and Li01 combination treatment	Fei et al., 2022.
23	Goji juice fermented by <i>Lactobacillus plantarum</i> , <i>Lactobacillus reuteri</i> and <i>Streptococcus thermophilus</i>	Probiotics-fermentation enhanced the anti-ulcerative colitis function of goji berry juice and modulated gut microbiota	Liu et al., 2021.
24	<i>Lactobacillus plantarum</i> CBT LP3 (KCTC 10782BP)	Effective anti-inflammatory effects, with increased induction of Treg and restoration of goblet cells, suppression of proinflammatory cytokines	Kim et al., 2020.
25	<i>Bifidobacterium breve</i> , CCFM683	Improved intestinal epithelial barriers, restored gut microbiota	Chen et al., 2020b.
26	<i>Bifidobacterium infantis</i> , <i>Lactobacillus acidophilus</i> , <i>Enterococcus faecalis</i> with (quadruple probiotics, Pqua) or without (triple probiotics, P-tri) aerobic <i>Bacillus cereus</i>	Effective (Aerobe-contained Piqua was a powerful adjuvant therapy for chronic colitis, via restoring the intestinal microflora and recovering the multi-barriers in the inflamed gut)	Chen et al., 2020c.
27	<i>Lactococcus lactis</i> subsp. <i>lactic</i> JCM5805	Effective (high-dose administration deteriorates	Komaki et al., 2020.

(Continued)

TABLE 2 Continued

Sr. no.	Probiotic Used	Outcome	Reference
		intestinal inflammation)	
28	<i>Lactobacillus bulgaricus</i>	Regulates the inflammatory response and prevents Colitis-associated cancer	Silveira et al., 2020.
29	<i>Lactobacillus</i> and <i>Bifidobacterium</i> species	Significantly induced remission in UC patients	Agraib et al., 2022.
30	<i>Bacillus coagulans</i> Unique IS-2	Showed beneficial effects when administered along with standard medical treatment	Bamola et al., 2022.
31.	<i>Bifid Triple Viable Capsules</i>	IL-1, TNF- α , and IL-10 had higher decreases in a test group	Cui et al., 2004.
32	<i>Lactobacillus casei</i> DG	Both orally and rectally given probiotics have shown SS improvement in clinical and histological scores	D'Inca et al., 2011.
33	<i>Bifidobacterium longum</i>	Sigmoidoscopy scores (SS) and blood-serological markers TNF- α and IL-1 were reduced. Both clinical activity index (CAI) and bowel habit index (BHI) were reduced in a test group	Furrie et al., 2005.
34	<i>Bifidobacterium breve</i>	The endoscopic score of the treatment group was significantly lower. Myeloperoxidase analysis (MPO) amounts in the lavage solution (LS) significantly decreased	Ishikawa et al., 2011.
35	<i>Bifid Triple Viable Capsules</i>	Higher decrease in UCDAI scores and symptoms in the test group. TNF- α and IL-8 were decreased in a test group	Huang et al., 2018.
36	<i>Enterococcus faecium</i> , <i>Lactobacillus plantarum</i> , <i>Streptococcus thermophilus</i> , <i>Bifidobacterium lactis</i> , <i>Lactobacillus acidophilus</i> , <i>Bifidobacterium longum</i>	SS differences in decrease of endoscopic and clinical index score. The test group achieved a higher decrease	Kamarli et al., 2019.
37	<i>Bifidobacterium breve</i> , <i>Bifidobacterium bifidum</i> , <i>Lactobacillus acidophilus</i> YIT 0168	CAI score, endoscopic score, and histological score were significantly lower in the treatment group	Kato et al., 2004.

(Continued)

TABLE 2 Continued

Sr. no.	Probiotic Used	Outcome	Reference
38	<i>E.coli</i> Nissle 1917 (Serotype O6: K5: H1)	No significant differences both in CAI scores and relapse rates. Relapse-free time differences were also NS	Kruis et al., 1997.
39	<i>E.coli</i> Nissle 1917 (Serotype O6: K5: H1)	NS differences in decrease of clinical symptoms and blood-serological markers between groups. Both groups had decreased inflammation markers and symptoms	Arribas et al., 2009.
40	<i>Bifidobacterium breve</i> , <i>Bifidobacterium bifidum</i> , <i>Lactobacillus acidophilus</i> YIT 0168	NS differences in both relapse-free survival and clinical deterioration	Matsuoka et al., 2018.
41	<i>E.coli</i> Nissle 1917 (Serotype O6: K5: H1)	The dose depended on efficacy in both remission time and endoscopic findings	Matthes et al., 2010.
42	<i>L. paracasei</i> , <i>L. plantarum</i> , <i>L. acidophilus</i> , <i>L. delbrueckii</i> subsp <i>bulgaricus</i> , <i>B. longum</i> , <i>B. breve</i> , <i>B. infantis</i> , <i>Streptococcus thermophilus</i>	More patients achieved remission in the test group	Ng et al., 2010.
43	<i>Lactobacillus salivarius</i> , <i>Lactobacillus acidophilus</i> , <i>Bifidobacterium bifidus</i> strain BGN4	The better improvement compared to the control	Palumbo et al., 2016.
44	<i>Bifidobacterium longum</i> BB536	Significant decrease in UCDAI scores and endoscopic index in a test group	Tamaki et al., 2015.
45	<i>L. acidophilus</i> strain LA-5 and <i>B. animalis</i> subsp. <i>lactis</i> strain BB-12	More patients in the test group achieved remission. Median relapse time was longer in a test group	Wildt et al., 2011.
46	<i>Bifid Triple Viable Capsules</i>	The observation group had significantly lower scores in CDAI and UCAI as well as recurrence rate	Fan et al., 2019.
47	<i>Lactobacillus</i> spp.	Reduced fecal calprotectin (FCAL) in UC patients. No differences in IBD-QOL scores and blood-serological markers	Yilmaz et al., 2019.
48	<i>Bifidobacterium</i> and <i>Lactobacillus</i> with sulfasalazine and prednisone vs. sulfasalazine	Level decrease of CRP, TNF- α and IL-10 in both groups, significantly lower in the study group ($p < 0.05$); significantly	Su et al., 2018.

(Continued)

TABLE 2 Continued

Sr. no.	Probiotic Used	Outcome	Reference
		higher treatment effect in the study group (p < 0.05); higher infection rate in the control group (p < 0.05)	
49	VSL#3	75% of patients remained in remission during the study period, with no side effects	Venturi et al., 1999.
50	<i>Eschericia coli</i> Nissle 1917 (1×1011cfu/day)	Prebiotics induce remission as effectively as mesalazine (standard treatment)	Rembacken et al., 1999.
51	VSL#3	VSL#3 resulted in combined induction of remission/response rate of 70%, with no adverse effects	Bibiloni et al., 2005.
52	<i>Saccharomyces boulardii</i>	71% of patients remained in remission	Guslandi et al., 2003.
53	Yakult (1×1010 cfu/day)	73% of patients treated with probiotics remained in remission, while only 10% of placebo	Ishikawa et al., 2003.

has indicated that *Bifidobacterium adolescentis* can utilise XOS, while, *Lactobacillus rhamnosus*, *L. plantarum* and *Lactococcus lactis* have been found to efficiently metabolise oat -glucooligosaccharides. According to Van Laere et al. (2000), the fermentation process of arabinose-XOS derived from wheat meal by *bifidobacteria* can be attributed to the existence of xylanolytic enzyme systems such as xylosidase and a limited number of arabinosidases. The presence of a -D-xylosidase derived from *Bifidobacterium breve* K-110, as well as arabinosidases originating from *B. adolescentis* DSM20083 (Van Laere et al., 2000) and *B. breve* (Shin et al., 2000), has been documented in the literature. Zeng et al. (2007) conducted a study on the species *B. bifidum*, *B. adolescentis*, and *B. infantis*, and reported that these species exclusively exhibited arabinosidase and xylosidase activity, while no activity of, glucuronidase, or acetyl xylan esterase and xylanase was observed. Gullon et al. (2008) studied the fermentation process of XOS from rice husks using probiotic bacteria. They found that *Bifidobacterium adolescentis* CECT 5781 showed significantly greater growth compared to *Breve* CECT 4839, *Bifidobacterium longum* CECT 4503 and *Infantis* CECT 4551 in the presence of XOS. This highlights the importance of industrial development in prebiotics for enhancing food quality and human health.

A variety of functional foods containing prebiotics have been utilised in the production of biscuits, candies, tabletop sweeteners, frozen yoghurt, and other similar products (Davani-Davari et al., 2019). In Japan, the Foods for Specified Health Uses (FOSHU) programme has incorporated various prebiotics, such as FOS, IMO,




FIGURE 2
Characteristics of prebiotic.

XOS, lacto-sucrose, and lactulose oligosaccharides, into specific food products. Additional examples of prebiotic foods include yoghurts, cereals, cakes; cereal bars biscuits high in dietary fibre, powdered beverages, pasta, sauces, bread, infant formula products and various fruit juices (Desai, 2008). Prebiotics are commonly associated with specific types of dietary fibre, such as inulin and FOS. However, they can also be found in various food sources, including cultivated plants and wild plants. Below are several examples of food plants that are abundant in prebiotics; categorised accordingly as cultivated and wild plants.

Cultivated plants

- Bananas: Bananas contain FOS, which is a type of prebiotic that is fermented by beneficial bacteria in the gut. FOS has been shown to advance gut health, decrease inflammation, and improve the immune system. (Sabater-Molina et al., 2009)
- Garlic: Garlic contains inulin, a prebiotic fibre that nourishes gut bacteria. It is also known for its antimicrobial properties. (Davani-Davari et al., 2019)
- Legumes: Legumes, such as beans, lentils, and peas, are excellent sources of resistant starch, a type of prebiotic that is fermented by beneficial bacteria in the gut. Resistant starch has been shown to increase gut health, decrease the risk of chronic diseases, and benefit weight management. (Mirmiran, 2014)
- Oats: Oats are a great source of beta-glucans, a category of soluble fibre that is fermented by beneficial bacteria in the gut. Beta-glucans has been shown to improve gut health, reduce cholesterol levels, and help with weight management. (Sabater-Molina et al., 2009)
- Whole grains: Whole grains, such as brown rice, quinoa, and barley, are excellent sources of resistant starch and other prebiotics. (Sabater-Molina et al., 2009)
- Onion: Like garlic, onions are rich in inulin and can promote the growth of helpful gut bacteria. (Davani-Davari et al., 2019)
- *Asparagus*: *Asparagus* is a good source of inulin, a type of soluble fibre that is fermented by beneficial bacteria in the gut. Inulin has been shown to improve gut health, boost immunity, and decrease the risk of chronic disorders. (Sabater-Molina et al., 2009)
- *Jerusalem artichoke*: *Jerusalem artichokes* are high in inulin content, making them an excellent prebiotic food. (Sabater-Molina et al., 2009)
- Chicory Root: Chicory root is often used as a source of inulin in supplements, which is proven to increase the growth of *Bifidobacterium*. It can be consumed raw or roasted (Davani-Davari et al., 2019).

Wild Plants:

- *Taraxacum officinale* (Dandelion greens): A very common plant, dandelion grows wild almost everywhere. Dandelion is a native of Europe. In India, it is found in the Himalayas.

Nutritionally, the dandelion has remarkable value. Dandelion greens are wild plant that contains inulin and other prebiotic fibres, along with various nutrients (Wirngo et al., 2016).

- *Arctium lappa L* (Burdock Root): Burdock root is a wild plant that contains inulin and has been traditionally used as a medicinal food with prebiotic properties. This species is native to the temperate regions of the Old World, from Scandinavia to the Mediterranean, and from the British Isles through Russia, and the Middle East to India, China, Taiwan and Japan. (Moro and Clerici, 2021).
- *Cichorium intybus L* (Chicory Greens): Wild chicory greens are rich in inulin and can be consumed in salads or cooked as a side dish. The Chicory crop is cultivated in a few States, mostly Uttar Pradesh, and Gujarat. These two states account for 97% of the total production of Chicory in India (Nwafor et al., 2017).
- *Musa paradisiaca* (Plantain): Plantain leaves are a common wild plant that contains prebiotic fibres, among other beneficial compounds. Plantain fruit is widely consumed in Nigeria, Africa and some other parts of the world (Ukwah et al., 2014).
- *Urtica Dioica* (Nettles): Nettles are known for their numerous health benefits and can provide prebiotic effects due to their fibre content. It is most common in Europe, North America, North Africa, and parts of Asia (Behzad et al., 2018).

These are just a few examples, and there are many other cultivated and wild plants that can offer prebiotic benefits. Combining a type of these prebiotic-rich foods into your diet can aid in promoting a healthy gut microbiome (Crittenden and Playne, 2008; Davani-Davari et al., 2019). For a product (food or supplement) to be believed to be a prebiotic, it must meet the below-mentioned conditions (Gibson et al., 2017).

Prebiotic enhance the proliferation and metabolic function of selected bacterial strains that exhibit beneficial impacts on overall well-being in the following ways.

- Lower the pH of the substances presents in the intestines.
- To exhibit resistance to hydrolysis and the effects of gastrointestinal enzymes.
- To avoid being absorbed in the upper gastrointestinal tract.
- To maintain a suitable environment for one or more beneficial microorganisms within the colon.
- To maintain stability throughout the food processing procedure,

7.1 Prebiotics in UC

Prebiotics, mostly fermentable carbohydrates, promote local or systemic health (Pandey et al., 2015; Akram et al., 2019). Prebiotics can alter the intestinal microbiota, improve the intestinal barrier, and

promote the growth of beneficial microbes in the digestive tract, which produce host-beneficial metabolites. (Aljuraiban et al., 2023) Like probiotics, clinical trials on prebiotics in specific diseases are difficult to prove. Thus, prebiotic data in UC patients is carefully regulated. Prebiotics may help treat UC by supplementing fermentable carbohydrate-containing fibre fractions. Promoting specific bacteria and/or their metabolites achieves this (Pandey et al., 2015; Rasmussen and Hamaker, 2017). Prebiotics may help maintain remission or low clinical disease activity. Prebiotics used in UC studies are oligosaccharides and inulin (Rasmussen and Hamaker, 2017). The mechanism for the ameliorative effect of prebiotics in UC studies is illustrated in Figure 3. Prebiotics encourage the growth of beneficial microorganisms, which compete with harmful species and produce beneficial fermentation substances like SCFAs, which have immunomodulatory properties and influence toll-like receptor-4 signalling and pro-inflammatory cytokines. (Van der Beek et al., 2017)

Casellas et al. (2007) studied the effects of mesalazine treatment, oligofructose-enriched inulin, and placebo on patients with mild to moderate UC. Oral oligofructose-enriched inulin was well-received and reduced faecal calprotectin levels. Faecal calprotectin has been employed as a reliable and quantifiable indicator of intestinal inflammation, with a significant correlation observed between its levels and the assessment of disease activity in UC, as determined through histologic and endoscopic methods (Konikoff and Denson, 2006). In Japan, germinated barley food (GBF) has been explored as a therapeutic intervention for UC. GBF, a dietary fibre and protein-rich in glutamine, has been shown to reduce clinical activity in individuals with UC and is an effective treatment option for maintaining

remission. The safety profile of the treatment is noteworthy, as the study conducted by Hanai et al. (2004) found no incidence of side effects in the study group associated with the use of GBF. Nevertheless, further clinical trials are required to provide additional confirmation regarding the efficacy of dietary fibre as a prebiotic in the management of UC. Prebiotics has been extensively utilised in the management and treatment of UC. Table 3 presents a comprehensive compilation of various studies that have examined the advantageous effects of prebiotics in the management of UC.

8 Synbiotics

Gibson’s proposition of prebiotics was accompanied by a rationale for the potential advancements that could be achieved through the combination of prebiotics with probiotics, leading to the establishment of what he referred to as Synbiotics (De Vrese and Schrezenmeir, 2008). The term “synbiotics” pertains to the amalgamation of prebiotics and probiotics to improve the well-being of humans or animals (Markowiak and Śliżewska, 2017). According to research conducted by Perrin et al. (2001) and Sharma and Shukla (2016), the probiotic bacteria found in synbiotic food products make selective use of prebiotics as a substrate for their growth. A group of specialists from the International Scientific Association for Probiotics and Prebiotics got together and rethought the concept of synbiotics. Synbiotic treatment aims to improve the endurance and metabolic function of beneficial probiotic strains in the gut microbiota. Common combinations include lactobacilli and *bifidobacteria* with oligosaccharides, inulin,




TABLE 3 Studies of prebiotics in the management of UC.

Sr. No.	Prebiotic used	Outcome Results	Reference
1	Plantago ovata seed	Maintain remission as effectively as mesalamine, Increased in fecal butyrate.	Fernandez-Banares, 1999.
2	Synergy 1 (inulin and oligofructose) 6 g/day and Bifidobacterium longum	Reduction of defensins 2, 3, and 4, TNF - α and IL- 1	Furrie et al., 2005.
3	Germinated barley foodstuff (30 g/day)	Reduction of clinical activity index scores and increase in stool butyrate concentrations.	Mitsuyama et al., 1998.
4	Polysaccharide from <i>Scutellaria baicalensis</i> Georgi	Effective (attenuated body weight loss, reduced DAI, ameliorated colonic pathological damage, and decreased MPO activity)	Cui et al., 2021.
5	Synthetic glycans	Synthetic glycans increase survival, reduce weight loss, and improve clinical scores in mouse models of colitis	Tolonen et al., 2022.
6	GOS, FOS along with FMT	Treatment with FMT plus a prebiotic blend restores the structure of the intestinal flora and increased the levels of acetic acid, butyric acid, FFAR3, and ZO-1	Qian et al., 2022.
7	Alpha D-glucan from marine fungus <i>Phoma herbarum</i> YS4108	Effective (significantly increased butyrate, isovaleric acid levels, and prominent alterations on specific microbiota)	Liu et al., 2016.
8	Stachyose	Increased beneficial microbiota and bacterial diversity to alleviate acute colitis in mice	He et al., 2020.
9	<i>Dictyophora indusiate</i> polysaccharide	Effective (modulates gut microbiota)	Kanwal et al., 2020.
10	FMG or dealcoholised muscadine wine	Effective (reduced dysbiosis in the colon)	Li et al., 2020.
11	GFO	Effective (prevented and attenuated colitis symptoms and GI dysmotility, reducing populations of harmful bacteria and increasing SCFAs)	K-da et al., 2020.
12	β -fructans (oligofructose and inulin)	Did not prevent symptomatic relapses in UC patients but reduced the severity of biochemical relapse and increased anti-inflammatory metabolites	Valcheva et al., 2022.
13	Oral microencapsulated sodium butyrate (BLM)	BLM supplementation appears to be a valid add-on therapy to maintain remission in patients with UC	Vernero et al., 2020.

(Continued)

TABLE 3 Continued

Sr. No.	Prebiotic used	Outcome Results	Reference
14	Oligofructose-enriched inulin	Less significant (scFOS on rectal sensitivity may require higher doses and may depend on the subgroup)	Valcheva et al., 2018.

or fibres as prebiotics. This approach reduces systemic inflammation by increasing the population of bacteria producing SCFAs and providing substrates for fermentation (Pandey et al., 2015).

According to their classification, synbiotics can be categorised into two distinct classes: complementary and synergistic. A complementary synbiotic is comprised of a probiotic and a prebiotic, which collectively provide one or more health benefits without requiring interdependent functions. According to Swanson et al. (2020), a synergistic synbiotic is composed of a substrate that is specifically utilised by microorganisms that are co-administered. These guidelines are expected to be efficacious in establishing expectations regarding the comprehension of the interplay between pre- and probiotics as well as the advancement of synbiotic products for the promotion of health and therapeutic interventions. Numerous reports suggest that the consumption of synbiotic foods has a beneficial impact on the health and nutritional status of the host. The study conducted by Yang et al. (2005) revealed that the administration of synbiotics had a dual effect on the faecal sample. Firstly, it was observed that synbiotics increased the abundance of probiotic bacteria, specifically *Lactobacillus* and *Bifidobacterium*, while simultaneously reducing the presence of coliform bacteria. Additionally, the test group exhibited an improvement in the levels of various digestive enzymes, including lactase, sucrase, lipase and isomaltase. Recently, there has been a report indicating that the consumption of synbiotics has significantly decreased various cardiovascular risk factors, the prevalence of metabolic syndrome, and markers of insulin resistance among elderly individuals (Cicero et al., 2021). The ability of *bifidobacteria* to metabolise prebiotics is dependent on the species, which is a valuable characteristic for modulating the gut microbiota through the use of specialised prebiotics (Bielecka et al., 2002; Biedrzycka and Bielecka, 2004). The β -fructofuranosidase enzyme derived from *Bifidobacterium adolescentis* G1 exhibits a preference for fructooligomers over inulin as its substrate. This preference is also observed in *B. bifidum*. In contrast, it has been observed that *B. longum* and *B. animalis* possess the ability to hydrolyze a diverse array of FOS and XOS, including those derived from inulin (Bruno et al., 2002). According to a study conducted by Bruno et al. (2002), food products contain the greatest concentration of viable *bifidobacteria*. Likewise, the efficacy of *B. longum* supplemented with FOS is greater in curd. According to Martinez-Villaluenga and Gomez (2007), the probiotic *B. lactis* possesses the enzymes β -glucosidase and β -fructofuranosidase, which enable it to metabolise oligosaccharides present in fermented milk. This metabolic activity promotes the growth and metabolism of the probiotic organism. The therapeutic potentials and health

benefits of different probiotics, prebiotics, and synbiotics are presented in Table 4.

8.1 Synbiotics in ulcerative colitis

Synbiotics are products that combine probiotics and prebiotics, resulting in a synergistic interaction. This concept was introduced

to address challenges in probiotic survival, especially during transit through the upper gastrointestinal tract. The use of a synbiotic enhances probiotic colonisation efficacy and facilitates the proliferation of probiotic strains (Pandey et al., 2015).

There is a limited body of research examining the impact of a synbiotic supplement on individuals diagnosed with UC. Table 5 presents a compilation of the studies that have been cited most frequently and illustrated in Figure 4. Wong et al. (2022)

TABLE 4 Therapeutic potential and health benefits of probiotics, prebiotics and synbiotics.

Biotic types	Sources	Diseases	Health effects	Mechanism of action	References
Probiotics	<i>Lactobacillus acidophilus</i> and <i>Bifidobacterium infantis</i>	Intestinal infections	Inhibition of <i>Staphylococcus aureus</i> , <i>Salmonella typhimurium</i> , <i>Yersinia enterocolitica</i> , <i>Clostridium perfringens</i> and <i>Aeromonas hydrophila</i>	Production of organic acids, bacteriocins and other primary metabolites, such as hydrogen peroxide, carbon dioxide and diacetyl	Shahani and Chandan, 1979; Laroia and Martin, 1990; Mishra and Lambert, 1996; Van der Meer and Bovee-Oudenhoven, 1998
	<i>L. casei</i> , <i>L. acidophilus</i> and <i>B. bifidum</i>	Immune enhancement	Data not available	Enhancement in non-specific (e.g. phagocyte function, NK cell activity) and specific (e.g. antibody and cytokine production) host immune responses	Kaila et al., 1992; Schiffrin et al., 1995; Gill, 1998
	<i>L. acidophilus</i> , <i>S. thermophilus</i> , <i>B. longum</i> , <i>L. rhamnosus</i> GG and <i>B. bifidum</i>	Diarrhoeal infections	Inhibitions of <i>Escherichia coli</i> , <i>Salmonella</i> , <i>Shigella</i> , <i>Clostridium difficile</i> and rotavirus	Production of organic acids, bacteriocins, hydrogen peroxide, carbon dioxide and diacetyl	Merson et al., 1976; Barefoot and Klaenhammer, 1983; Tojo et al., 1987; Oksanen et al., 1990; Siitonen et al., 1990; Hilton et al., 1997.
	<i>B. longum</i> , <i>L. casei</i> Shirota, <i>L. acidophilus</i> , <i>Bifidobacterium</i> spp. and <i>L. rhamnosus</i> GG	Cancer	Inhibition of tumour formation and proliferation	Inhibition of carcinogens and procarcinogens, bacteria converting procarcinogens to carcinogens, immune system activation, and reduced faecal enzyme levels	Lidbeck et al., 1991; Reddy and Rivenson, 1993; Goldin et al., 1996; McIntosh, 1996.
	<i>L. acidophilus</i>	Hypercholesterolaemia	Reduction of cholesterol levels	Assimilation of cholesterol and deconjugation of bile salts	Gilliland and Speck, 1977; Buck and Gilliland, 1994.
	<i>L. acidophilus</i> , <i>B. angulatum</i> , <i>B. breve</i> , <i>B. bifidum</i> and <i>B. longum</i>	Lactose intolerance	Utilisation of lactose	Production of β-D-galactosidase which hydrolyzes lactose	Kilara and Shahani, 1976; Hughes and Hoover, 1995.
	<i>L. acidophilus</i> and <i>Bifidobacterium</i> spp. Production of	Reduction of peptic ulcer, gastro-oesophageal reflux, non-ulcer dyspepsia and gastric cancer	Inhibition of <i>Helicobacter pylori</i>	Lactic and acetic acids, bacteriocins etc	Berrada et al., 1991; Lambert and Hull, 1996; Lankaputhra et al., 1996.
	<i>L. rhamnosus</i> GG	Food allergy	Help to relieve intestinal inflammation and hypersensitivity reactions in infants with food allergies	Hydrolyse the complex casein to smaller peptides and amino acids and hence decrease the proliferation of mitogen-induced human lymphocytes	(Sutas et al., 1996; Majamaa and Isolauri, 1997)
Prebiotics	Inulin from chicory roots	-	-	Stimulate the growth of <i>Bifidobacterium</i>	Gibson et al., 1995.
	Neosugar	-	-	Metabolised by the resident microbes in the colon including <i>bifidobacteria</i> , <i>Enterococcus faecalis</i> , <i>E. faecium</i> , <i>Bacteroides vulgates</i> , etc	Desai, 2008; Guarino et al., 2020.
	Isomalto-oligosaccharides (IMO)	-	Local and systemic Th-1-like immune response and	<i>Bifidobacterium</i> and the <i>Bacteroides</i> groups can utilise IMO	Kohmoto et al., 1988; Wang et al., 2014.

(Continued)

TABLE 4 Continued

Biotic types	Sources	Diseases	Health effects	Mechanism of action	References
	from miso, soy sauce and honey		regulation of immune function, balancing the dysbiosis of gut microbiota		
	Xylooligosaccharides (XOS) from fruits, bamboo shoots, vegetables, honey, etc.	-	-	<i>B. adolescentis</i> utilizes xylobiose and xylotriose, whereas <i>L. lactis</i> , <i>L. rhamnosus</i> and <i>L. plantarum</i> utilise oat β -glucosaccharides	Okazaki et al., 1990.
Synbiotics	Food products containing <i>B. animalis</i> and amylose cornstarch	-	-	Promote the growth of <i>bifidobacteria</i>	Bruno et al., 2002.
	Curd containing <i>B. longum</i> and fructooligosaccharide (FOS)	-	Decrease cardiovascular risk factors, metabolic syndrome prevalence and markers of insulin resistance in elderly patients	Promote the growth of <i>B. longum</i>	Hughes and Hoover, 1995; Linares et al., 2017; Cicero et al., 2021.
	Oral synbiotic preparation containing <i>L. plantarum</i> and FOS	Sepsis in early infancy	Significant reduction in sepsis and lower respiratory tract infections	Promotes growth of <i>L. plantarum</i> ATCC202195	Panigrahi et al., 2017.
	Synbiotics containing five probiotics (<i>L. plantarum</i> , <i>L. delbrueckii</i> spp. <i>bulgaricus</i> , <i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>Bifidobacterium bifidum</i>) and inulin	-	-	Adult subjects with NASH (non-alcoholic <i>steatohepatitis</i>) demonstrated a significant reduction of IHTG (intrahepatic triacylglycerol)	Wong et al., 2013.
	Synbiotic products containing <i>L. rhamnosus</i> , <i>Bifidobacterium lactis</i> , inulin and oligofructose	Hepatic conditions	-	Increased level of intestinal IgA, reduced blood cholesterol levels and lower blood pressure	Pathmakanthan et al., 2002; Perez-Conesa et al., 2006.
	<i>L. rhamnosus</i> CGMCC1.3724 and inulin	Obesity	Weight loss	Reduction in leptin increase in Lachnospiraceae	Sanchez et al., 2014.
	<i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>B. bifidum</i> , <i>B. longum</i> , <i>E. faecium</i> and FOS	Obesity	Changes in anthropometric measurements	Decrease in TC, LDL-C and total oxidative stress serum levels	Ipar et al., 2015.
	<i>L. sporogenes</i> and inulin	Type 2 diabetes	-	Significant reduction in serum insulin levels and homeostatic model assessment cell function	Tajadadi-Ebrahimi et al., 2014.
	<i>L. casei</i> , <i>L. rhamnosus</i> , <i>S. thermophilus</i> , <i>B. breve</i> , <i>L. acidophilus</i> , <i>B. longum</i> , <i>L. bulgaricus</i> and FOS	Insulin resistance syndrome	The levels of fasting blood sugar and insulin resistance improved significantly	-	Eslamparast et al., 2014.
	<i>L. rhamnosus</i> GG, <i>B. lactis</i> Bb12 and inulin	Cancer	Increase in probiotics in stools and decrease in <i>Clostridium perfringens</i> led to increase in the IL2 in polypectomised patients	Increased production of interferon- γ	Safavi et al., 2013.

investigated the impact of prebiotic mixtures, probiotic mixtures, and synbiotics on colitis in a murine model. The findings of the study indicated that the administration of synbiotic treatment had a protective effect on the structure of the colon, as evidenced by the preservation of its integrity. Furthermore, there was an observed

increase in the expression of occludin, a protein involved in maintaining the tight junctions between cells, which is indicative of improved barrier function. Additionally, the treatment was found to be effective in reducing the infiltration of cells into the colon and resulted in alterations to the gut microbiome, enhancements in

TABLE 5 Summary of the most relevant studies involving synbiotics treatments in UC.

Sr. No.	Synbiotic	Outcome Results	Reference
1	<i>Bifidobacterium longum</i> plus inulin-oligofructose Treatment time: one month	Sigmoidoscopy scores ↓ β defensins 2, 3, and 4 ↓ CRP TNF-α ↓; IL-1β ↓; IgA and IgG production ↔	Furrie, 2005.
2	<i>Bifidobacterium longum</i> plus <i>psyllium</i> Treatment time: 4 weeks	CRP ↓ IBDQ (total, bowel, systemic, emotional, and social functional scores) ↑	Fujimori et al., 2009.
3	<i>Lactobacillus Paracasei</i> B 20160 + XOS Treatment time: 8 weeks	Serum IL-6, IL-8 ↓ Serum TNF-α, IL-1-β ↔ PBMC, IL-8 ↓	Federico et al. (2009)
4	<i>Bifidobacterium breve</i> strain Yakult plus galactooligosaccharides Treatment time: one-year	MPO ↓ Bacteroidaceae ↓ fecal pH ↓ clinical status: improved	Ishikawa et al., 2011.
5	<i>Lactobacillus acidophilus</i> LA-5®, <i>Lactobacillus delbrueckii</i> subsp. <i>bulgaricus</i> LBY-27, <i>Bifidobacterium animalis</i> subsp. <i>lactis</i> BB-12® and <i>Streptococcus thermophilus</i> STY-31™ plus oligofructose Treatment time: One month	Microflora spectrum ↔	Ahmed et al., 2013.
6	<i>Streptococcus faecalis</i> T-110 JPC, <i>Clostridium butyricum</i> TO-A, <i>Bacillus mesentericus</i> TO-A JPC, <i>Lactobacillus sporogenes</i> plus prebiotic Treatment time: 3 months	Severity score ↓ Steroid intake ↓ Relapse during follow-up (3 months) ↓ Duration of remission ↑	Malathi et al., 2019.
7	A symbiotic which concluded six probiotics: <i>Enterococcus faecium</i> , <i>Lactobacillus plantarum</i> , <i>Streptococcus thermophilus</i> , <i>Bifidobacterium lactis</i> , <i>Lactobacillus acidophilus</i> , <i>Bifidobacterium longum</i> and fructooligosaccharide	The change in the CRP and sedimentation values had a statistically significant decrease in the synbiotic group (P = 0.003). The improvement in clinical activity was significantly higher in the synbiotic group (p < 0.05)	Kamarli et al., 2019.
8	<i>Lactobacillus plantarum</i> LP90 and Soluble dietary fibre obtained from <i>Lentinula edodes</i> by-products	Alleviated colitis	Xue et al., 2023.
9	<i>Bifidobacterium infantis</i> and <i>Bifidobacterium</i>	Increased diversity of the microbiome and be	Ivanovska et al., 2017.

(Continued)

TABLE 5 Continued

Sr. No.	Synbiotic	Outcome Results	Reference
	<i>longum</i> and Equal parts FOS, GOS and XOS	associated with more SCFAs, and less gut inflammation	
10	<i>L. paracasei</i> and <i>Opuntia humifusa</i> extract (mucilage + pectin)	Effective (greater abundance of <i>L. paracasei</i> in fecal microbial analysis, lower serum corticosterone levels, lower TNF-α levels in the colon tissue	Seong et al., 2020.
11	VSL#3 and Yacon (6% FOS + inulin)	Preservation of intestinal architecture, improve intestinal integrity, increased expression of antioxidant enzymes and concentration of organic acids	Dos Santos Cruz et al., 2020.
12	<i>Lactobacillus acidophilus</i> , <i>L. Rhamnosus</i> , <i>Bifidobacterium lactis</i> and Inulin	Increased the proportion of helpful bacteria and regulated the balance of intestinal microbiota, reduced the degree of inflammation in acute colitis mice	Wang et al., 2019.
13	LGG and Tagatose	Effective (gut microbiota composition recovered from the dysbiosis caused by DSS treatment)	Son et al., 2019.
14	<i>L. Casey</i> 01 and Oligofructose-enriched inulin	Assessment of colonic damage, inflammation scoring, MPO and microbiological studies are done and its effects on the UC	Ivanovska et al., 2017.
15	<i>Lactobacillus casei</i> , <i>Lactobacillus acidophilus</i> , <i>Lactobacillus rhamnosus</i> , <i>Lactobacillus bulgaricus</i> , <i>Bifidobacterium breve</i> , <i>Bifidobacterium longum</i> , <i>Streptococcus thermophiles</i> and FOS	Mitigated symptoms in patients with UC and suggested to use pre probiotics in the standard treatment, particularly in those with more than five years of the disease	Amiriani et al., 2020.
16	Food products containing <i>B. animalis</i> and amylose corn starch Data not available Data not available	<i>L. plantarum</i> utilise oat β-glucooligosaccharides to Promote the growth of <i>bifidobacteria</i>	Bruno et al., 2002.
17	Curd containing <i>B. longum</i> and fructooligosaccharide (FOS)	Promote the growth of <i>B. longum</i>	Hughes and Hoover, 1995; Linares et al., 2017; Cicero et al., 2021

(Continued)

TABLE 5 Continued

Sr. No.	Synbiotic	Outcome Results	Reference
18	Oral synbiotic preparation containing <i>L. plantarum</i> and FOS	Promotes growth of <i>L. plantarum</i> ATCC202195	Panigrahi et al., 2017.
19	Synbiotics containing five probiotics (<i>L. plantarum</i> , <i>L. delbrueckii</i> spp. <i>bulgaricus</i> , <i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>Bifidobacterium bifidum</i>) and inulin	Adult subjects with NASH (non-alcoholic steatohepatitis) demonstrated a significant reduction of IHTG (intrahepatic triacylglycerol)	Wong et al., 2013.
20	Synbiotic product containing <i>L. rhamnosus</i> , <i>Bifidobacterium lactis</i> , inulin and oligofructose	Increased level of intestinal IgA, reduced blood cholesterol levels and lower blood pressure	Pathmakanthan et al., 2002; Perez-Conesa et al., 2006.
21	<i>L. rhamnosus</i> CGMCC1.3724 and inulin	Reduction in leptin increase in Lachnospiraceae	Sanchez et al., 2014.
22	<i>L. acidophilus</i> , <i>L. rhamnosus</i> , <i>B. bifidum</i> , <i>B. longum</i> , <i>E. faecium</i> and FOS	Decrease in TC, LDL-C and total oxidative stress serum levels	Ipar et al., 2015.
23	<i>L. sporogenes</i> and inulin	Significant reduction in serum insulin levels and homeostatic model assessment cell function	Tajadadi-Ebrahimi et al., 2014.
24	<i>L. casei</i> , <i>L. rhamnosus</i> , <i>S. thermophilus</i> , <i>B. breve</i> , <i>L. acidophilus</i> , <i>B. longum</i> , <i>L. bulgaricus</i> and FOS	The levels of fasting blood sugar and insulin resistance improved significantly	Eslamparast et al., 2014.
25	<i>L. plantarum</i> La-5, <i>B. animalis</i> subsp. <i>lactis</i> BB-12 and dietary fibres	Improvement in the IBS score and satisfaction in bowel movement reported	Smid et al., 2016.
26	<i>L. rhamnosus</i> GG, <i>B. lactis</i> Bb12 and inulin	Increased production of interferon- γ	Safavi et al., 2013.

Increase, ↑; decrease, ↓; equivalent, ↔.

colonic integrity, and the suppression of markers associated with inflammation. Although the existing studies have yielded promising results thus far, it is important to note that the methodologies employed often exhibit inconsistencies, lack proper description, and/or suffer from inadequate study design. Additionally, it is worth mentioning that the majority of cases have been characterised by a notably limited number of registered patients.

9 Safety and considerations in ulcerative colitis treatment

When it comes to treating ulcerative colitis, ensuring safety and considering various factors are of paramount importance. Here are some key safety considerations to keep in mind:

9.1 Medical supervision

Ulcerative colitis treatment should always be conducted under the guidance and supervision of a qualified healthcare professional. They will assess the severity of the condition, consider individual patient factors, and prescribe appropriate medications or therapies (Kerner et al., 2014).

9.2 Monitoring and follow-up

Regular monitoring is crucial to evaluating the effectiveness of the treatment and assessing any potential side effects or complications. Patients should attend follow-up appointments as scheduled and communicate any changes in symptoms or concerns to their healthcare provider (Click and Regueiro, 2019).

9.3 Medication safety

Patients need to follow the prescribed medication regimen diligently. They should be aware of potential side effects and know when to seek medical attention. Certain medications used in ulcerative colitis treatment, such as immunosuppressants or biologics, may require additional precautions and monitoring due to their impact on the immune system (Click and Regueiro, 2019).

9.4 Individualised approach


Each patient with ulcerative colitis is exceptional, and management plans should be designed to meet their special needs. Factors such as age, disease severity, comorbidities, and medication tolerance should be taken into consideration when developing a personalised treatment strategy (Kerner et al., 2014).

9.5 Lifestyle modifications

In addition to medication, lifestyle changes can play a major role in managing ulcerative colitis. These may include dietary changes, stress management techniques, regular exercise, and getting sufficient rest. Patients should work with healthcare professionals, such as dietitians and psychologists, to make appropriate lifestyle adjustments (Click and Regueiro, 2019).

9.6 Risk-benefit assessment

Every treatment decision involves a careful evaluation of the potential risks and benefits. Patients and healthcare providers should discuss the expected outcomes, potential side effects, and long-term implications of various treatment options. This shared decision-making process ensures that the chosen treatment aligns with the patient's goals and preferences (Kerner et al., 2014).



9.7 Complementary and alternative therapies

If considering complementary or alternative therapies, patients should consult with their healthcare provider to ensure safety and assess potential interactions with conventional treatments. It is important to be cautious of unproven or unsupported therapies that may promise a cure without scientific evidence (Click and Regueiro, 2019).

9.8 Patient education and support

Patient education about their condition, treatment options, and self-care is essential. Ulcerative colitis patients can benefit from educational resources and support groups. In conclusion, ulcerative colitis treatment must prioritise safety and individual factors. Patients can manage their condition while minimising risks and maximising treatment benefits with medical supervision, regular monitoring, personalised approaches, and informed decision-making (Kerner et al., 2014).

10 Future perspectives and research

UC significantly impacts individuals' health and quality of life, with treatment options including aminosalicylates, corticosteroids, immunomodulators, biological therapies, and surgical interventions. However, some patients have not responded well to these treatments, leading to prolonged suffering and reduced quality of life. Alternative therapies like experimental drugs are considered, but their effectiveness remains uncertain and comes with risks and complications. As we gain more understanding of the pathophysiology of UC and its relationship with the microbiome, there will be a rise in research trials to determine the efficacy of emerging treatments. Scholars are currently investigating "next-generation" probiotics, such as *Clostridium clusters* IV, XIVa, and XVIII, *Bacteroides uniformis*, *Bacteroides fragilis*, *Akkermansia*

muciniphila, *Eubacterium hallii*, and *Faecalibacterium prausnitzii* as potential alternatives.

Collaborative engagement between patients and healthcare providers is crucial for determining the most suitable treatment approach. Emerging treatments like prebiotics, probiotics, and synbiotics have gained attention for their potential therapeutic benefits in improving gut health and treating various conditions. For patients with IBD, traditional medications may not always work for every patient, and their side effects can be severe. In clinical practice, incorporating probiotics, prebiotics, and synbiotics holds great promise for transforming the approach to managing ulcerative colitis, offering patients a safer and more focused alternative to conventional therapeutic interventions. This will facilitate our understanding of the interplay between human physiological processes and the microbiome.

11 Discussion

The search for a definitive solution for UC continues to be a persistent effort, with existing therapeutic interventions primarily focused on maintaining a state of remission. In the management of this condition, conventional therapeutic approaches encompass the utilisation of aminosalicylates, corticosteroids, immunomodulators, and biological therapies. These interventions are employed with the objectives of attaining remission, averting relapses, and facilitating the restoration of mucosal integrity. Current therapeutic approaches encompass the use of monoclonal antibodies that specifically target proinflammatory cytokines, adhesion molecules, and T-cell activation, as well as anti-inflammatory cytokines like IL10 and TGF-. However, the administration of these medications can lead to various adverse effects, both at the systemic and local levels. These effects may include headaches, nausea, abdominal pain, loss of appetite, vomiting, and the development of a rash. The use of corticosteroids for an extended period is discouraged due to the potential increase in susceptibility to infections (Mowat et al., 2011). In contemporary times, functional food is increasingly

recognised for its dual capacity to ensure nutritional security and confer health benefits on the consumer (Mijan and Lim, 2018). Promising advancements have been observed in the utilisation of probiotics, prebiotics, and synbiotics for the management of UC. Bacterial strains, such as *Lactobacillus* and *Bifidobacterium* species, have been linked to beneficial outcomes concerning symptom management and the maintenance of remission in individuals with a diagnosis of UC. By encouraging the growth of advantageous microorganisms and producing anti-inflammatory fatty acids, prebiotics supports gut health and digestion. Non-digestible fibres in foods like whole grains, garlic, and bananas support gut flora, lowering the risk of gastrointestinal conditions like irritable bowel syndrome and enhancing immune function. Prebiotic-rich foods are an excellent way to support digestive health and uphold general wellbeing through diet. The synergistic effects of prebiotics and probiotics allow for the effective management of UC (Fujimori et al., 2009). Therefore, it is imperative to conduct a comprehensive examination of these interventions to effectively mitigate any potential negative consequences and develop strategies for long-term supervision. The management of UC can be tailored to the individual by combining conventional treatments with probiotics, prebiotics, and synbiotics. The evaluated combination therapies demonstrate encouraging synergistic effects through the simultaneous targeting of various aspects of disease pathogenesis, including inflammation, immune dysregulation, and gut microbiota imbalance. Nevertheless, it is of utmost importance to conduct a thorough assessment of treatment interactions, contraindications, and individual patient variables to guarantee the safety and effectiveness of the intervention. It is imperative to undertake additional research, employing meticulously crafted clinical trials, to ascertain the most advantageous strains, compositions, and therapeutic regimens for probiotics, prebiotics, and synbiotics within the particular framework of UC. The aforementioned advancements provide empirical support for the capacity to enhance therapeutic outcomes, alleviate adverse effects, and establish innovative and well-tolerated strategies for managing UC. The current literature on probiotics and prebiotics in IBD is influenced by significant heterogeneity, with varying study designs, doses, and outcomes. The study populations varied, with some focusing on active disease patients and others on remission maintenance. Most studies enrolled small numbers of patients, limiting statistical power. The exact mechanisms of action of probiotics, prebiotics, and synbiotics are not yet fully understood. Insufficient evidence on probiotic dosages and immunological mechanisms is needed to establish health claims. The interaction between microbiota, host, and prebiotic components is also limited. Clinical trials and validation studies with larger sample sizes require understanding of these interactions. Limited published literature in manufacturing processes and formulation further needs improvement. (Astó et al., 2019)

References

Aachary, A. A., and Prapulla, S. G. (2010). Xylooligosaccharides (XOS) as an emerging prebiotic: Microbial synthesis, utilization, structural characterization,

In conclusion, the utilisation of probiotics, prebiotics, and synbiotics in the management of UC showcases a promising advancement with potential implications. These interventions present innovative strategies for manipulating the gut microbiota, reducing inflammation, and improving overall health. Additional investigation is required to determine the most effective utilisation and potential incorporation of these interventions into individualised therapeutic approaches for managing ulcerative colitis. Through the adoption of innovative approaches in the treatment of UC and the consideration of individual patient requirements, we have the potential to advance towards a more optimistic future for individuals afflicted by this complex inflammatory bowel disease.

Author contributions

AJ: Conceptualization, Data curation, Writing – original draft. SJ: Conceptualization, Supervision, Validation, Writing – review & editing. SV: Writing – review & editing. AS: Writing – review & editing. BK: Writing – review & editing.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Acknowledgments

The authors are thankful to all the authorities of Bharati Vidyapeeth (Deemed to be a University) for their overall support and encouragement.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

bioactive properties, and applications. *Compr. Rev. Food Sci. Food Saf.* 10, 2–16. doi: 10.1111/j.1541-4337.2010.00135.x

- Agraib, L. M., Yamani, M. I., Tayyem, R., Abu-Sneineh, A. T., and Rayyan, Y. M. (2022). Probiotic supplementation induces remission and changes in the immunoglobulins and inflammatory response in active ulcerative colitis patients: A pilot, randomized, double-blind, placebo-controlled study. *Clin. Nutr. ESPEN* 51, 83–91. doi: 10.1016/j.clnesp.2022.08.020
- Ahmed, J., Reddy, B. S., Molbak, L., Leser, T. D., and MacFie, J. (2013). Impact of probiotics on colonic microflora in patients with colitis: A prospective double blind randomised crossover study. *Int. J. Surg.* 11, 1131–1136. doi: 10.1016/j.ijvs.2013.08.019
- Akram, W., Garud, N., and Joshi, R. (2019). Role of inulin as prebiotics on inflammatory bowel disease. *Drug Discoveries Ther.* 13, 1–8. doi: 10.5582/dtd.2019.01000
- Alatab, S., Sepanlou, S. G., Ikuta, K., Vahedi, H., Bisignano, C., Safiri, S., et al. (2020). The global, regional, and national burden of inflammatory bowel disease in 195 countries and territories 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Gastroenterol. Hepatol.* 5, 17–30. doi: 10.1016/s2468-1253(19)30333-4
- Aljuraiban, G. S., Alghasani, S. S., Sabico, S., Alshammari, S., Aljazairy, E. A., and Al-Musharaf, S. (2023). Types of fiber and gut microbiota composition and diversity among arab females. *Saudi J. Biol. Sci.* 30, 103767. doi: 10.1016/j.sjbs.2023.103767
- Amiriani, T., Rajabli, N., Faghani, M., Besharat, S., Roshandel, G., Akhavan, T., et al. (2020). Effect of Lactocare[®] synbiotic on disease severity in ulcerative colitis: A randomized placebo-controlled double-blind clinical trial. *Middle East J. Digestive Dis.* 12, 27–33. doi: 10.15171/mejdd.2020.160
- Ananthakrishnan, A. N. (2015). Epidemiology and risk factors for IBD. *Nat. Rev. Gastroenterol. Hepatol.* 12, 205–217. doi: 10.1038/nrgastro.2015.34
- Arribas, B., Rodriguez, C. M., Camuesco, D., Comalada, M., Bailon, E., Utrilla, P., et al. (2009). A probiotic strain of *Escherichia coli*, Nissle 1917, given orally exerts local and systemic anti-inflammatory effects in lipopolysaccharide-induced sepsis in mice. *Br. J. Pharmacol.* 157, 1024–1033. doi: 10.1111/j.1476-5381.2009.00270.x
- Astó, E., Méndez, I., Audivert, S., Farrán-Codina, A., and Espadaler, J. (2019). The efficacy of probiotics, prebiotic inulin-type fructans, and synbiotics in human ulcerative colitis: A systematic review and meta-analysis. *Nutrients* (11), 293. doi: 10.3390/nu11020293
- Bamola, V. D., Dubey, D., Samanta, P., Kedia, S., Madempudi, R. S., Neelamraju, J., et al. (2022). Effect of *Bacillus coagulans* unique is-2 in inflammatory bowel disease (IBD): A randomized controlled trial. *medRxiv* 78, 1–29. doi: 10.1101/2021.07.18.21260556
- Barcenilla, C., Ducic, M., Lopez, M., Prieto, M., and Alvarez-Ordóñez, A. (2021). Application of lactic acid bacteria for the biopreservation of meat products: A systematic review. *Meat Sci.* 183, 1–13.108661. doi: 10.1016/j.meatsci.2021.108661
- Barefoot, S. F., and Klaenhammer, T. R. (1983). Detection and activity of lactacin B, a bacteriocin produced by *Lactobacillus acidophilus*. *Appl. Environ. Microbiol.* 45, 1808–1815. doi: 10.1128/aem.45.6.1808-1815.1983
- Barnaba, V., and Sinigaglia, F. (1997). Molecular mimicry and T cell-mediated autoimmune disease. *J. Exp. Med.* 185, 1529–1532. doi: 10.1084/jem.185.9.1529
- Basso, P. J., Camara, N. O., and Sales-Campos, H. (2019). Microbial-based therapies in the treatment of inflammatory bowel disease – an overview of human studies. *Front. Pharmacol.* 9. doi: 10.3389/fphar.2018.01571
- Behzad, K., Asghar, K., Mohammad, A. K., and Rezvan, P. (2018). Probiotic yogurt formulated with nettle (*Urtica dioica*) extract, a compound with dual functionalities: *Bifidobacterium* growth promoter and *Helicobacter pylori* growth inhibitor. *BioMed. J. Sci. Tech Res.* 10, 7532–7538. doi: 10.26717/BJSTR.2018.10.001888
- Berrada, N., Lemeland, J.-F., Laroche, G., Thouvenot, P., and Piaia, M. (1991). *Bifidobacterium* from fermented milks: Survival during Gastric transit. *J. Dairy Sci.* 74, 409–413. doi: 10.3168/jds.s0022-0302(91)78183-6
- Bibiloni, R., Fedorak, R. N., Tannock, G. W., Madsen, K. L., Gionchetti, P., Campieri, M., et al. (2005). VSL3 probiotic-mixture induces remission in patients with active ulcerative colitis. *Am. J. Gastroenterol.* 100, 1539–1546. doi: 10.1111/j.1572-0241.2005.41794.x
- Biedrzycka, E., and Bielecka, M. (2004). Prebiotic effectiveness of fructans of different degrees of polymerization. *Trends Food Sci. Technol.* 15, 170–175. doi: 10.1016/j.tifs.2003.09.014
- Bielecka, M., Biedrzycka, E., and Majkowska, A. (2002). Selection of probiotics and prebiotics for synbiotics and confirmation of their *in vivo* effectiveness. *Food Res. Int.* 35, 125–131. doi: 10.1016/s0963-9969(01)00173-9
- Bischoff, S. C., Escher, J., Hebuterne, X., Klęk, S., Krznaric, Z., Schneider, S., et al. (2020). Espen practical guideline: Clinical nutrition in inflammatory bowel disease. *Clin. Nutr.* 39, 632–653. doi: 10.1016/j.clnu.2019.11.002
- Biarnason, I., Sission, G., and Hayee, B. (2019). A randomised, double-blind, placebo-controlled trial of a multi-strain probiotic in patients with asymptomatic ulcerative colitis and crohn's disease. *Inflammopharmacology* 27, 465–473. doi: 10.1007/s10787-019-00595-4
- Britton, R. A., Hoffmann, D. E., and Khoruts, A. (2021). Probiotics and the microbiome—how can we help patients make sense of probiotics? *Gastroenterology* 160, 614–623. doi: 10.1053/j.gastro.2020.11.047
- Bruno, F. A., Lankaputhra, W. E. V., and Shah, N. P. (2002). Growth, viability and activity of *Bifidobacterium* spp. in skim milk containing prebiotics. *J. Food Sci.* 67, 2740–2744. doi: 10.1111/j.1365-2621.2002.tb08807.x
- Buck, L. M., and Gilliland, S. E. (1994). Comparisons of freshly isolated strains of *Lactobacillus acidophilus* of human intestinal origin for ability to assimilate cholesterol during growth. *J. Dairy Sci.* 77, 2925–2933. doi: 10.3168/jds.s0022-0302(94)77233-7
- Casellas, F., Borrueal, N., Torrejon, A., Varela, E., Antolin, M., Guarner, F., et al. (2007). Oral oligofructose-enriched inulin supplementation in acute ulcerative colitis is well tolerated and associated with lowered faecal calprotectin. *Alimentary Pharmacol. Ther.* 25, 1061–1067. doi: 10.1111/j.1365-2036.2007.03288.x
- Çavdar, M., and Çavdar, M. (2023). Alternative therapeutic applications used in the treatment of ulcerative colitis: probiotics, prebiotics, synbiotics and fecal microbiota transplantation. *Prog. Nutr.* doi: 10.23751/pn.v25i2.13775
- Chapman, C. M., Gibson, G. R. O., and Rowland, I. (2011). Health benefits of probiotics: Are mixtures more effective than single strains? *Eur. J. Nutr.* 50, 1–17. doi: 10.1007/s00394-010-0166-z
- Chen, Y., Zhang, L., Hong, G., Huang, C., Qian, W., Bai, T., et al. (2020c). Probiotic mixtures with aerobic constituent promoted the recovery of multi-barriers in DSS-induced chronic colitis. *Life Sciences* 240, 117089. doi: 10.1016/j.lfs.2019.117089
- Chen, Y., Jin, Y., Stanton, C., Ross, R. P., Wang, Z., Zhao, J., et al. (2020b). Dose-response efficacy and mechanisms of orally administered cla-producing *Bifidobacterium breve* CCFM683 on DSS-induced colitis in mice. *J. Funct. Foods* 75, 104245. doi: 10.1016/j.jff.2020.104245
- Chen, X., Liu, Y., Jin, J., Liu, H., Hao, Y., Zhang, H., et al. (2021). YbfA regulates the sensitivity of *Escherichia coli* K12 to *Plantaricin* BM-1 via the BasS/BasR two-component regulatory system. *Front. Microbiol.* 12. doi: 10.3389/fmicb.2021.659198
- Chen, P., Xu, H., Tang, H., Zhao, F., Yang, C., Kwok, L. Y., et al. (2020a). Modulation of gut mucosal microbiota as a mechanism of probiotics-based adjunctive therapy for ulcerative colitis. *Microbial Biotechnol.* 13, 2032–2043. doi: 10.1111/1751-7915.13661
- Cicero, A. F. G., Fogacci, F., Bove, M., Giovannini, M., and Borghi, C. (2021). Impact of a short-term synbiotic supplementation on metabolic syndrome and systemic inflammation in elderly patients: a randomized placebo-controlled clinical trial. *Eur. J. Nutr.* 60, 655–663. doi: 10.1007/s00394-020-02271-8
- Click, B., and Regueiro, M. (2019). A practical guide to the safety and monitoring of new IBD therapies. *Inflammatory Bowel Dis.* 25, 831–842. doi: 10.1093/ibd/izy313
- Cosovanu, C., and Neumann, C. (2020). The many functions of Foxp3+ Regulatory T cells in the intestine. *Front. Immunol.* 11. doi: 10.3389/fimmu.2020.600973
- Crittenden, R., and Playne, M. J. (2008). “Nutrition News. Facts and functions of prebiotics, probiotics and synbiotics,” in *Handbook of Probiotics and Prebiotics*. Eds. Y. K. Lee and S. Salminen (Hoboken, NJ, USA; Manhattan, KS, USA: Wiley-Interscience, Kansas State University), 2008 535–582.
- Cui, H. H., Chen, C. L., Wang, J. D., Yang, Y. J., Cun, Y., Wu, J. B., et al. (2004). Effects of probiotic on intestinal mucosa of patients with ulcerative colitis. *World journal of gastroenterology* 10, 1521–1525. doi: 10.3748/wjg.v10.i10.1521
- Cui, L., Guan, X., Ding, W., Luo, Y., Wang, W., Bu, W., et al. (2021). *Scutellaria baicalensis* Georgi polysaccharide ameliorates DSS-induced ulcerative colitis by improving intestinal barrier function and modulating gut microbiota. *Int. J. Biol. Macromolecules* 166, 1035–1045. doi: 10.1016/j.ijbiomac.2020.10.259
- Dai, M., Li, Y., Xu, L., Wu, D., Zhou, Q., Li, P., et al. (2021). A novel bacteriocin from *Lactobacillus pentosus* ZFM94 and its antibacterial mode of action. *Front. Nutr.* 8. doi: 10.3389/fnut.2021.710862
- Dalile, B., Van Oudenhove, L., Vervliet, B., and Verbeke, K. (2019). The role of short-chain fatty acids in microbiota–gut–brain communication. *Nat. Rev. Gastroenterol. Hepatol.* 16, 461–478. doi: 10.1038/s41575-019-0157-3
- Davani-Davari, D., Negahdaripour, M., Karimzadeh, I., Seifan, M., Mohkam, M., Masoumi, S. J., et al. (2019). Prebiotics: definition, types, sources, mechanisms, and clinical applications. *Foods (Basel Switzerland)* 8, 92. doi: 10.3390/foods8030092
- Derikx, L., Dieleman, L. A., and Hoentjen, F. (2016). Probiotics and prebiotics in ulcerative colitis. *Best Pract. Res. Clin. Gastroenterology* 30, 55–71. doi: 10.1016/j.bpg.2016.02.005
- Derwa, Y., Gracie, D. J., Hamlin, P. J., and Ford, A. C. (2017). Systematic review with meta-analysis: the efficacy of probiotics in inflammatory bowel disease. *Alimentary Pharmacol. Ther.* 46, 389–400. doi: 10.1111/apt.14203
- Desai, A. R. (2008). *Strain identification, viability and probiotics properties of Lactobacillus casei*. Ph.D. thesis, School of Biomedical and Health Sciences (Werribee Campus Victoria, Australia: Victoria University).
- De Vrese, M., and Schrezenmeir, J. (2008). Probiotics, prebiotics, and synbiotics. *Food Biotechnol.* 111, 1–66. doi: 10.1007/10_2008_097
- D'Inca, R., Barollo, M., Scarpa, M., Grillo, A. R., Brun, P., Vettorato, M. G., et al. (2010). Rectal administration of *Lactobacillus casei* DG modifies flora composition and Toll-Like Receptor expression in colonic mucosa of patients with mild ulcerative colitis. *Digestive Dis. Sci.* 56, 1178–1187. doi: 10.1007/s10620-010-1384-1
- Dominguez-Bello, M. G., Costello, E. K., Contreras, M., Magris, M., Hidalgo, G., Fierer, N., et al. (2010). Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc. Natl. Acad. Sci.* 107, 11971–11975. doi: 10.1073/pnas.1002601107
- Dos Santos Cruz, B. C., da Silva Duarte, V., Giacomini, A., Corich, V., Oliveira de Paula, S., da Silva, F. L., et al. (2020). Synbiotic VSL3 and Yacon-based product modulate the intestinal microbiota and prevent the development of pre-neoplastic

- lesions in a colorectal carcinogenesis model. *Appl. Microbiol. Biotechnol.* 104, 8837–8857. doi: 10.1007/s00253-020-10863-x
- Eslamparast, T., Zamani, F., Hekmatdoost, A., Sharafkhan, M., Egtesad, S., Malekzadeh, R., et al. (2014). Effects of synbiotic supplementation on insulin resistance in subjects with the metabolic syndrome: a randomised, double-blind, placebo-controlled pilot study. *Br. J. Nutr.* 112, 438–445. doi: 10.1017/S0007114514000919
- Fan, H., Du, J., Liu, X., Zheng, W. W., Zhuang, Z. H., Wang, C. D., et al. (2019). Effects of pentasa-combined probiotics on the microflora structure and prognosis of patients with inflammatory bowel disease. *Turkish J. Gastroenterol.* 30, 680–685. doi: 10.5152/tjg.2019.18426
- Fathizadeh, H., Pakdel, F., Saffari, M., Esmaceli, D., Heravi, M. M., Dao, S., et al. (2022). Bacteriocins: recent advances in its application as an antimicrobial alternative. *Curr. Pharm. Biotechnol.* 23, 1028–1040. doi: 10.2174/1389201022666210907121254
- Federico, A., Tuccillo, C., Grossi, E., Abbiati, R., Garbagna, N., Romano, M., et al. (2009). The effect of a new symbiotic formulation on plasma levels and peripheral blood mononuclear cell expression of some pro-inflammatory cytokines in patients with ulcerative colitis: a pilot study. *Eur. Rev. Med. Pharmacol. Sci.* 13, 285–293. doi: 10.1200/JCO.2008.21.3991
- Fei, Y., Zhang, S., Han, S., Qiu, B., Lu, Y., Huang, W., et al. (2022). The role of dihydroresveratrol in enhancing the synergistic effect of *Ligilactobacillus salivarius* Li01 and resveratrol in ameliorating colitis in mice. *Res. (Washington D.C.)* 2022, 9863845. doi: 10.34133/2022/9863845
- Fernandez-Banares, F. (1999). Randomized clinical trial of *Plantago ovata* seeds (dietary fiber) as compared with mesalazine in maintaining remission in ulcerative colitis. *Am. J. Gastroenterol.* 94, 427–433. doi: 10.1016/s0002-9270(98)00753-9
- Fujimori, S., Gudis, K., Mitsui, K., Seo, T., Yonezawa, M., Tanaka, S., et al. (2009). A randomized controlled trial on the efficacy of synbiotic versus probiotic or prebiotic treatment to improve the quality of life in patients with ulcerative colitis. *Nutr. (Burbank Los Angeles County Calif.)* 25, 520–525. doi: 10.1016/j.nut.2008.11.017
- Fuller, R. (1989). Probiotics in man and animals. *J. Appl. Bacteriol.* 66, 365–378.
- Furrie, E., Macfarlane, S., Kennedy, A., Cummings, J. H., Walsh, S. V., O'neil, D. A., et al. (2005). Synbiotic therapy (*Bifidobacterium Longum*/synergy 1) initiates resolution of inflammation in patients with active ulcerative colitis: A randomized controlled pilot trial. *Gut* 54, 242–249. doi: 10.1136/gut.2004.044834
- Fuss, I. J., and Strober, W. (2008). The role of IL-13 and NK T cells in experimental and human ulcerative colitis. *Mucosal Immunol.* 1, S31–S33. doi: 10.1038/mi.2008.40
- Gibson, G. R., Beatty, E. R., Wang, X., and Cummings, J. H. (1995). Selective stimulation of *bifidobacteria* in the human colon by oligofructose and inulin. *Gastroenterology* 108, 975–982. doi: 10.1016/0016-5085(95)90192-2
- Gibson, G. R., Hutkins, R., Sanders, M. E., Prescott, S. L., Reimer, R. A., Salminen, S. J., et al. (2017). Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of probiotics. *Nat. Rev. Gastroenterol. Hepatol.* 14, 491–502. doi: 10.1038/nrgastro.2017.75
- Gibson, G. R., and Roberfroid, M. B. (1995). Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics. *J. Nutr.* 125, 1401–1412. doi: 10.1093/jn/125.6.1401
- Gill, H. (1998). Stimulation of the immune system by lactic cultures. *Int. Dairy J.* 8, 535–544. doi: 10.1016/s0958-6946(98)00074-0
- Gilliland, S. E., and Speck, M. L. (1977). Deconjugation of bile acids by intestinal lactobacilli. *Appl. Environ. Microbiol.* 33, 15–18. doi: 10.1128/aem.33.1.15-18.1977
- Goldin, B. R., Gualtieri, L. J., and Moore, R. P. (1996). The effect of *Lactobacillus* GG on the initiation and promotion of DMH induced intestinal tumors in the rat. *Nutr. Cancer* 25, 197–204. doi: 10.1080/01635589609514442
- Guarino, M., Altomare, A., Emerenziani, S., Di Rosa, C., Ribolsi, M., Balestrieri, P., et al. (2020). Mechanisms of action of probiotics and their effects on gastro-intestinal disorders in adults. *Nutrients* 12, 1037. doi: 10.3390/nu12041037
- Gullon, P., Moura, P., Esteves, M. P., Girio, F. M., Dominguez, H., and Parajo, J. C. (2008). Assessment on the fermentability of xylooligosaccharides from rice husks by probiotic bacteria. *J. Agric. Food Chem.* 56, 7482–7487. doi: 10.1021/jf800715b
- Guslandi, M., Giollo, P., and Testoni, P. A. (2003). A pilot trial of *Saccharomyces boulardii* in ulcerative colitis. *Eur. J. Gastroenterol. Hepatol.* 15, 697–698. doi: 10.1097/00042737-200306000-00017
- Hanai, H., Kanauchi, O., Mitsuyama, K., Andoh, A., Takeuchi, K., Takayuki, I., et al. (2004). Germinated barley foodstuff prolongs remission in patients with ulcerative colitis. *Int. J. Mol. Med.* 3, 643–647. doi: 10.3892/ijmm.13.5.643
- Hart, A. L., Al-Hassi, H. O., Rigby, R. J., Bell, S. J., Emmanuel, A. V., Knight, S. C., et al. (2005). Characteristics of intestinal dendritic cells in inflammatory bowel diseases. *Gastroenterology* 129, 50–65. doi: 10.1053/j.gastro.2005.05.013
- He, Q., He, L., Zhang, F., Jian, Z., Sun, J., Chen, J., et al. (2020). Stachyose modulates gut microbiota and alleviates dextran sulfate sodium-induced acute colitis in mice. *Saudi J. Gastroenterol.* 26, 153. doi: 10.4103/sjg.sjg_580_19
- Heller, F., Florian, P., Bojarski, C., Richter, J., Christ, M., Hillenbrand, B., et al. (2005). Interleukin-13 the key effector Th2 cytokine in ulcerative colitis that affects epithelial tight junctions, apoptosis, and cell restitution. *Gastroenterology* 129, 550–564. doi: 10.1016/j.gastro.2005.05.002
- Heller, F., Fromm, A., Gitter, A. H., Mankertz, J., and Schulzke, J. D. (2008). Epithelial apoptosis is a prominent feature of the epithelial barrier disturbance in intestinal inflammation: effect of pro-inflammatory interleukin-13 on epithelial cell function. *Mucosal Immunol.* 1, s58–s61. doi: 10.1038/mi.2008.46
- Henker, J., Laass, M., Baretton, G., Fischer, R., and Aust, D. (2008). Fallstricke bei der Zöliakiediagnostik. *Z. Für Gastroenterologie* 46, 675–680. doi: 10.1055/s-2008-1027409
- Higuchi, L. M., Khalili, H., Chan, A. T., Richter, J. M., Bousvaros, A., and Fuchs, C. S. (2012). A prospective study of cigarette smoking and the risk of inflammatory bowel disease in women. *Am. J. Gastroenterol.* 107, 1399–1406. doi: 10.1038/ajg.2012.196
- Hill, C., Guarner, F., Reid, G., Gibson, G., Merenstein, D., Pot, B., et al. (2014). The international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat. Rev. Gastroenterol. Hepatol.* 11, 506–514. doi: 10.1038/nrgastro.2014.66
- Hilton, E., Kolakowski, P., Singer, C., and Smith, M. (1997). Efficacy of *Lactobacillus* GG as a diarrheal preventive in travelers. *J. Travel Med.* 4, 41–43. doi: 10.1111/j.1708-8305.1997.tb00772.x
- Hu, K. A., and Gubatan, J. (2023). Gut microbiome-based therapeutics in inflammatory bowel disease. *Clin. Trans. Discovery* 3, 1–9. doi: 10.1002/ctd2.182
- Huang, M., Chen, Z., Lang, C., Chen, J., Yang, B., Xue, L., et al. (2018). Efficacy of mesalazine in combination with bifid triple viable capsules on ulcerative colitis and the resultant effect on the inflammatory factors. *Pakistan J. Pharm. Sci.* 31, 2891–2895.
- Huda-Faujan, N., Abdulmir, A., Fatimah, A., Anas, O. M., Shuhaimi, M., Yazid, A., et al. (2010). The impact of the level of the intestinal short chain fatty acids in inflammatory bowel disease patients versus healthy subjects. *Open Biochem. J.* 4, 53–58. doi: 10.2174/1874091x01004010053
- Hughes, D. B., and Hoover, D. G. (1995). Viability and enzymatic activity of *bifidobacteria* in milk. *J. Dairy Sci.* 78, 268–276. doi: 10.3168/jds.s0022-0302(95)76634-6
- Im, J. P., Ye, B. D., Kim, Y. S., and Kim, J. S. (2018). Changing treatment paradigms for the management of inflammatory bowel disease. *Korean J. Internal Med.* 33, 28–35. doi: 10.3904/kjim.2017.400
- Ipar, N., Aydogdu, S. D., Yildirim, G. K., Inal, M., Gies, I., Vandenplas, Y., et al. (2015). Effects of synbiotic on anthropometry, lipid profile and oxidative stress in obese children. *Beneficial Microbes* 6, 775–781. doi: 10.3920/bm2015.0011
- Ishikawa, H., Akedo, I., Umetsaki, Y., Tanaka, R., Imaoka, A., and Otani, T. (2003). Randomized controlled trial of the effect of *bifidobacteria*-fermented milk on ulcerative colitis. *J. Am. Coll. Nutr.* 22, 56–63. doi: 10.1080/07315724.2003.10719276
- Ishikawa, H., Matsumoto, S., Ohashi, Y., Imaoka, A., Setoyama, H., Umetsaki, Y., et al. (2011). Beneficial effects of probiotic *Bifidobacterium* and galacto-oligosaccharide in patients with ulcerative colitis: A randomized controlled study. *Digestion* 84, 128–133. doi: 10.1159/000322977
- Ivanovska, T. P., Mladenovska, K., Zhivikj, Z., Pavlova, M. J., Gjurovski, I., Ristoski, T., et al. (2017). Synbiotic loaded chitosan-Ca-alginate microparticles reduces inflammation in the TNBS model of rat colitis. *Int. J. Pharmaceutics* 527, 126–134. doi: 10.1016/j.ijpharm.2017.05.049
- Jakubczyk, D., Leszczynska, K., and Gorska, S. (2020). The effectiveness of probiotics in the treatment of inflammatory bowel disease (IBD)—A critical review. *Nutrients* 12, 1973. doi: 10.3390/nu12071973
- Jin, W., Ai, H., Huang, Q., Li, C., He, X., Jin, Z., et al. (2023). Preclinical evidence of probiotics in ulcerative colitis: a systematic review and network meta-analysis. *Front. Pharmacol.* 14. doi: 10.3389/fphar.2023.1187911
- Judge, T. A., and Lichtenstein, G. R. (2001). Refractory inflammatory bowel disease. *Curr. Treat Options Gastroenterol.* 4, 267–273. doi: 10.1007/s11938-001-0039-x
- Kaila, M., Isolauri, E., Soppi, E., Virtanen, E., Laine, S., and Arvilommi, H. (1992). Enhancement of the circulating antibody secreting cell response in human diarrhea by a human *Lactobacillus* strain. *Pediatr. Res.* 32, 141–144. doi: 10.1203/00006450-199208000-00002
- Kamarli, A. H., Akal, Y. E., and Akin, M. (2019). Effects of synbiotic therapy in mild-to-moderately active ulcerative colitis: A randomized placebo-controlled study. *Turkish J. Gastroenterol.* 30, 313–320. doi: 10.5152/tjg.2019.18356
- Kanwal, S., Joseph, T. P., Aliya, S., Song, S., Saleem, M. Z., Nisar, M. A., et al. (2020). Attenuation of DSS induced colitis by *Dictyophora indusiata* polysaccharide (DIP) via modulation of gut microbiota and inflammatory related signaling pathways. *J. Funct. Foods* 64, 103641. doi: 10.1016/j.jff.2019.103641
- Kato, K., Mizuno, S., Umetsaki, Y., Ishii, Y., Sugitani, M., Imaoka, A., et al. (2004). Randomized placebo-controlled trial assessing the effect of *bifidobacteria*-fermented milk on active ulcerative colitis. *Alimentary Pharmacol. Ther.* 20, 1133–1141. doi: 10.1111/j.1365-2036.2004.02268.x
- K-da, S., Peerakietkhajorn, S., Siringoringo, B., Muangnil, P., Wichienchot, S., and Khuituan, P. (2020). Oligosaccharides from *Gracilaria fisheri* ameliorate gastrointestinal dysmotility and gut dysbiosis in colitis mice. *J. Funct. Foods* 71, 104021. doi: 10.1016/j.jff.2020.104021
- Kerner, C., Lewis, J. D., and Osterman, M. T. (2014). “Safety considerations in the medical therapy of ulcerative colitis,” in *Medical Therapy of Ulcerative Colitis*. Ed. G. Lichtenstein (New York, NY: Springer).
- Khan, K. J., Ullman, T. A., Ford, A. C., Abreu, M. T., Abadir, A., Marshall, J. K., et al. (2011). Antibiotic therapy in inflammatory bowel disease: A systematic review and meta-analysis. *Am. J. Gastroenterology* 106, 661–673. doi: 10.1038/ajg.2011.72

- Khan, I., Wei, J., Li, A., Liu, Z., Yang, P., Jing, Y., et al. (2022). *Lactobacillus plantarum* strains attenuated DSS-induced colitis in mice by modulating the gut microbiota and immune response. *Int. Microbiol.* 25, 587–603. doi: 10.1007/s10123-022-00243-y
- Kilara, A., and Shahani, K. (1976). Lactase activity of cultured and acidified dairy products. *J. Dairy Sci.* 59, 2031–2035. doi: 10.3168/jds.s0022-0302(76)84484-0
- Kim, D. H., Kim, S., Ahn, J. B., Kim, J. H., Ma, H. W., Seo, D. H., et al. (2020). *Lactobacillus plantarum* CBT LP3 ameliorates colitis via modulating T cells in mice. *Int. J. Med. Microbiol.* 310, 151391. doi: 10.1016/j.ijmm.2020.151391
- Kmiec, Z., Cyman, M., and Ślebioda, T. J. (2017). Cells of the innate and adaptive immunity and their interactions in inflammatory bowel disease. *Adv. Med. Sci.* 62, 1–16. doi: 10.1016/j.advms.2016.09.001
- Kohmoto, T., Fukui, F., Takaku, H., Machida, Y., Arai, M., and Mitsuoka, T. (1988). Effect of isomaltoligosaccharides on human fecal flora. *bifidobacteria Microflora* 7, 61–69. doi: 10.12938/bifidus1982.7.2.61
- Komaki, S., Haque, A., Miyazaki, H., Matsumoto, T., and Nakamura, S. (2020). Unexpected effect of probiotics by *Lactococcus lactis* subsp. *lactis* against colitis induced by dextran sulfate sodium in mice. *J. Infection Chemotherapy* 26, 549–553. doi: 10.1016/j.jiac.2020.01.006
- Konikoff, M. R., and Denson, L. A. (2006). Role of fecal calprotectin as a biomarker of intestinal inflammation in inflammatory bowel disease. *Inflammatory Bowel Dis.* 12, 524–534. doi: 10.1097/00054725-200606000-00013
- Kruis, W., Fric, P., Pokrotnieks, J., Lukás, M., Fixa, B., Kascák, M., et al. (2004). Maintaining remission of ulcerative colitis with the probiotic *Escherichia coli* Nissle 1917 is as effective as with standard mesalazine. *Gut* 53, 1617–1623. doi: 10.1136/gut.2003.037747
- Kruis, W., Schütz, E., Fric, P., Fixa, B., Judmaier, G., and Stolte, M. (1997). Double-blind comparison of an oral *Escherichia coli* preparation and mesalazine in maintaining remission of ulcerative colitis. *Alimentary Pharmacol. Ther.* 11, 853–858. doi: 10.1046/j.1365-2036.1997.00225.x
- Kuhn, F., and Klar, E. (2015). Surgical principles in the treatment of ulcerative colitis. *Visceral Med.* 31, 246–250. doi: 10.1159/000438894
- Lambert, J., and Hull, R. (1996). Upper gastrointestinal tract disease and probiotics. *Asia Pacific J. Clin. Nutr.* 5, 31–35.
- Lankaputhra, W. E. V., Shah, N. P., and Britz, M. L. (1996). Evaluation of media for selective enumeration of *Lactobacillus acidophilus* and *Bifidobacterium* spp. *Food Aust.* 48, 113–118.
- Laroia, S., and Martin, J. (1990). *bifidobacteria* as possible dietary adjuncts in cultured dairy products—a review. *Cult Dairy Prod J.* 25, 18–22.
- Lavelle, A., and Sokol, H. (2020). Gut microbiota-derived metabolites as key actors in inflammatory bowel disease. *Nat. Rev. Gastroenterol. Hepatol.* 17, 223–237. doi: 10.1038/s41575-019-0258-z
- Li, P., Xiao, N., Zeng, L., Xiao, J., Huang, J., Xu, Y., et al. (2020). Structural characteristics of a mannoglycan isolated from Chinese yam and its treatment effects against gut microbiota dysbiosis and DSS-induced colitis in mice. *Carbohydr. Polymers* 250, 116958. doi: 10.1016/j.carbpol.2020.116958
- Lidbeck, A., Allinger, U. G., Orrhage, K. M., Ottova, L., Brismar, B., Gustafsson, J., et al. (1991). Impact of *Lactobacillus acidophilus* supplements on the faecal microflora and soluble faecal bile acids in colon cancer patients. *Microbial Ecol. Health Dis.* 4, 81–88. doi: 10.3109/08910609109140267
- Linares, D. M., Gomez, C., Renes, E., Fresno, J. M., Tornadijo, M. E., Ross, R. P., et al. (2017). Lactic acid bacteria and *bifidobacteria* with potential to design natural biofunctional health-promoting dairy foods. *Front. Microbiol.* 8. doi: 10.3389/fmicb.2017.00846
- Liu, Y., Fang, H., Liu, H., Cheng, H., Pan, L., Hu, M., et al. (2021). Goji berry juice fermented by probiotics attenuates dextran sodium sulfate-induced ulcerative colitis in mice. *J. Funct. Foods* 83, 104491. doi: 10.1016/j.jff.2021.104491
- Liu, R., Li, Y., and Zhang, B. (2016). The effects of konjac oligosaccharide on TNBS-induced colitis in rats. *Int. Immunopharmacol.* 40, 385–391. doi: 10.1016/j.intimp.2016.08.040
- Machiels, K., Joossens, M., Sabino, J., De Preter, V., Arijis, I., Eeckhaut, V., et al. (2013). A decrease of the butyrate-producing species *Roseburia hominis* and *Faecali bacterium prausnitzii* defines dysbiosis in patients with ulcerative colitis. *Gut* 63, 1275–1283. doi: 10.1136/gutjnl-2013-304833
- Mackowiak, P. A. (2013). Recycling Metchnikoff: Probiotics, the intestinal microbiome and the quest for long life. *Front. Public Health* 1. doi: 10.3389/fpubh.2013.00052
- Majamaa, H., and Isolauri, E. (1997). Probiotics: A novel approach in the management of food allergy. *J. Allergy Clin. Immunol.* 99, 179–185. doi: 10.1016/s0091-6749(97)70093-9
- Malathi, K., Nandini, R., Dhanasekar, K. R., and Shilpa, B. N. (2019). A randomized open label study to evaluate the efficacy and tolerability of synbiotic in the treatment of ulcerative colitis. *J. Vaccine Clin. Trials* 5, 1–4. doi: 10.35248/2475-3181.5.165
- Manichanh, C., Borrrel, N., Casellas, F., and Guarner, F. (2012). The gut microbiota in IBD. *Nat. Rev. Gastroenterol. Hepatol.* 9, 599–608. doi: 10.1038/nrgastro.2012.152
- Mariat, D., Firmesse, O., Levenez, F., Guimaraes, V. D., Sokol, H., Dore, J., et al. (2009). The Firmicutes/bacteroidetes ratio of the human microbiota changes with age. *BMC Microbiol.* 9, 1–6. doi: 10.1186/1471-2180-9-123
- Markowiak, P., and Ślizewska, K. (2017). Effects of probiotics, prebiotics, and synbiotics on human health. *Nutrients* 9, 1021. doi: 10.3390/nu9091021
- Martinez-Villaluenga, C., and Gomez, R. (2007). Characterization of *bifidobacteria* as starters in fermented milk containing raffinose family of oligosaccharides from lupin as prebiotic. *Int. Dairy J.* 17, 116–122. doi: 10.1016/j.idairyj.2006.02.003
- Matsuoka, K., Uemura, Y., Kanai, T., Kunisaki, R., Suzuki, Y., Yokoyama, K., et al. (2018). Efficacy of *Bifidobacterium breve* fermented milk in maintaining remission of ulcerative colitis. *Digestive Dis. Sci.* 63, 1910–1919. doi: 10.1007/s10620-018-4946-2
- Matthes, H., Krummenerl, T., Giensch, M., Wolff, C., and Schulze, J. (2010). Clinical trial: probiotic treatment of acute distal ulcerative colitis with rectally administered *Escherichia coli* Nissle 1917 (EcN). *BMC Complementary Altern. Med.* 10, 1–8. doi: 10.1186/1472-6882-10-13
- McIntosh, G. H. (1996). Probiotics and colon cancer prevention. *Asia Pacific J. Clin. Nutr.* 5, 48–52.
- Merson, M. H., Morris, G. K., Sack, D. A., Wells, J. G., Feeley, J. C., Sack, R. B., et al. (1976). Travelers' Diarrhea in Mexico. *New Engl. J. Med.* 294, 1299–1305. doi: 10.1056/nejm197606102942401
- Miele, E., Pascarella, F., Giannetti, E., Quaglietta, L., Baldassano, R. N., and Staiano, A. (2009). Effect of a probiotic preparation (VSL#3) on induction and maintenance of remission in children with ulcerative colitis. *Am. J. Gastroenterol.* 104, 437–443. doi: 10.1038/ajg.2008.118
- Miele, E., Shamir, R., Aloï, M., Assa, A., Braegger, C., Bronsky, J., et al. (2018). Nutrition in pediatric inflammatory bowel disease: A position paper on behalf of the porto inflammatory bowel disease group of the European Society of Pediatric Gastroenterology, Hepatology and Nutrition. *J. Pediatr. Gastroenterol. Nutr.* 66, 687–708. doi: 10.1097/mpg.0000000000001896
- Mijan, M. A., and Lim, B. O. (2018). Diets, functional foods, and nutraceuticals as alternative therapies for inflammatory bowel disease: Present status and future trends. *World J. Gastroenterol.* 24, 2673–2685. doi: 10.3748/wjg.v24.i25.2673
- Mirmiran, P. (2014). Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications: A review. *World J. Diabetes* 5, 267–281. doi: 10.4239/wjcd.v5.i3.26
- Mishra, C., and Lambert, J. (1996). Production of anti-microbial substances by probiotics. *Asia Pacific J. Clin. Nutr.* 5, 20–24.
- Mitsuyama, K., Saiki, T., Kanauchi, O., Iwanaga, T., Tomiyasu, N., Nishiyama, T., et al. (1998). Treatment of ulcerative colitis with germinated barley foodstuff feeding: a pilot study. *Alimentary Pharmacol. Ther.* 12, 1225–1230. doi: 10.1046/j.1365-2036.1998.00432.x
- Molodecky, N. A., Soon, I. S., Rabi, D. M., Ghali, W. A., Ferris, M., Chernoff, G., et al. (2012). Increasing incidence and prevalence of the inflammatory bowel diseases with time, based on systematic review. *Gastroenterology* 142, 46–54.e42. doi: 10.1053/j.gastro.2011.10.001
- Moro, T. M. A., and Clerici, T. P. S. (2021). Burdock (*Arctium lappa* L.) roots as a source of inulin-type fructans and other bioactive compounds: Current knowledge and future perspectives for food and non-food applications. *Food Res. Int. (Ottawa Ont.)* 141, 109889. doi: 10.1016/j.foodres.2020.109889
- Mowat, C., Cole, A., Windsor, A., Ahmad, T., Arnott, I., Driscoll, R., et al. (2011). Guidelines for the management of inflammatory bowel disease in adults. *Gut* 60, 571–607. doi: 10.1136/gut.2010.224154
- Nag, D., Farr, D., Raychaudhuri, S., and Withey, J. H. (2022). An adult zebrafish model for adherent-invasive *Escherichia coli* indicates protection from AIEC infection by probiotic *E. coli* Nissle. *iScience* 25, 104572. doi: 10.1016/j.isci.2022.104572
- Ng, S. C., Plamondon, S., Kamm, M. A., Hart, A. L., Al-Hassi, H. O., Guenther, T., et al. (2010). Immunosuppressive effects via human intestinal dendritic cells of probiotic bacteria and steroids in the treatment of acute ulcerative colitis. *Inflammatory Bowel Dis.* 16, 1286–1298. doi: 10.1002/ibd.21222
- Ng, S. C., Shi, H. Y., Hamidi, N., Underwood, F. E., Tang, W., Benchimol, E. I., et al. (2017). Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet* 390, 2769–2778. doi: 10.1016/s0140-6736(17)32448-0
- Nwafor, I. C., Shale, K., and Achilonu, M. C. (2017). Chemical composition and nutritive benefits of chicory (*Cichorium intybus*) as an ideal complementary and/or alternative livestock feed supplement. *Sci. World J.* 2017, 7343928. doi: 10.1155/2017/7343928
- Okazaki, M., Fujikawa, S., and Matsumoto, N. (1990). Effect of xylooligosaccharide on the growth of *bifidobacteria*. *Bifidobacteria Microflora* 9, 77–86. doi: 10.12938/bifidus1982.9.2.77
- Oksanen, P. J., Salminen, S., Saxelin, M., Hamalainen, P., Ihantola-Vormisto, A., Muurasiemi-Isoviita, L., et al. (1990). Prevention of travellers diarrhoea by *Lactobacillus* GG. *Ann. Med.* 22, 53–56. doi: 10.3109/07853899009147242
- Oliva, S., Di Nardo, G., Ferrari, F., Mallardo, S., Rossi, P., Patrizi, G., et al. (2011). Randomised clinical trial: the effectiveness of *Lactobacillus reuteri* ATCC 55730 rectal enema in children with active distal ulcerative colitis. *Alimentary Pharmacol. Ther.* 35, 327–334. doi: 10.1111/j.1365-2036.2011.04939.x
- Palumbo, V. D., Romeo, M., Gammazza, A. M., Carini, F., Damiani, P., Damiano, G., et al. (2016). The long-term effects of probiotics in the therapy of ulcerative colitis: A clinical study. *Biomed. Papers* 160, 372–377. doi: 10.5507/bp.2016.044

- Pandey, K. R., Naik, S. R., and Vakil, B. V. (2015). Probiotics, prebiotics and synbiotics- a review. *J. Food Sci. Technol.* 52, 7577–7587. doi: 10.1007/s13197-015-1921-1
- Panigrahi, P., Parida, S., Nanda, N. C., Satpathy, R., Pradhan, L., Chandel, D. S., et al. (2017). A randomized synbiotic trial to prevent sepsis among infants in rural India. *Nature* 548, 407–412. doi: 10.1038/nature23480
- Park, K. T., Ehrlich, O. G., Allen, J. I., Meadows, P., Szigethy, E. M., Henrichsen, K., et al. (2020). Corrigendum to the cost of Inflammatory Bowel Disease: An initiative from the Crohn's & Colitis Foundation. *Inflammatory Bowel Dis.* 26, 1118–1118. doi: 10.1093/ibd/izaa081
- Pathmakanthan, S., Walsh, M., and Bengmark, S. (2002). *Efficacy and tolerability treating acute distal ulcerative colitis with synbiotic enemas: a pilot trial (abstract)* (Geneva: United European Gastroenterology Week), 190.
- Perez-Conesa, D., Lopez, G., Abellan, P., and Ros, G. (2006). Bioavailability of calcium, magnesium and phosphorus in rats fed probiotic, prebiotic and synbiotic powder follow-up infant formulas and their effect on physiological and nutritional parameters. *J. Sci. Food Agric.* 86, 2327–2336. doi: 10.1002/jsfa.2618
- Perrin, S., Warchol, M., Grill, J., and Schneider, F. (2001). Fermentations of fructooligosaccharides and their components by *Bifidobacterium infantis* ATCC 15697 on batch culture in semi-synthetic medium. *J. Appl. Microbiol.* 90, 859–865. doi: 10.1046/j.1365-2672.2001.01317.x
- Petersen, A. M., Mirsepassi, H., Halkjær, S. I., Mortensen, E. M., Nordgaard-Lassen, I., and Kroghfelt, K. A. (2014). Ciprofloxacin and probiotic *Escherichia coli* Nissle add-on treatment in active ulcerative colitis: A double-blind randomized placebo-controlled clinical trial. *J. Crohn's Colitis* 8, 1498–1505. doi: 10.1016/j.crohns.2014.06.001
- Poggi, A., Benelli, R., Vene, R., Costa, D., Ferrari, N., Tosetti, F., et al. (2019). Human gut-associated natural killer cells in health and disease. *Front. Immunol.* 10. doi: 10.3389/fimmu.2019.00961
- Qian, X., Jiang, H., Wu, Y., Shao, M. H., He, J. W., Bao, X., et al. (2022). Fecal microbiota transplantation combined with probiotics ameliorates ulcerative colitis in mice. *Res. Square* 2022, 1–20. doi: 10.21203/rs.3.rs-1742266/v1
- Rasmussen, H. E., and Hamaker, B. R. (2017). Prebiotics and inflammatory bowel disease. *Gastroenterol. Clinics North America* 46, 783–795. doi: 10.1016/j.gtc.2017.08.004
- Reddy, B. S., and Rivenson, A. (1993). Inhibitory effect of *Bifidobacterium longum* on colon, mammary, and liver carcinogenesis induced by 2-amino-3-methylimidazo [4,5-f] quinoline, a food mutagen. *Cancer Res.* 53, 3914–3918.
- Rembacken, B., Snelling, A., Hawkey, P., Chalmers, D., and Axon, A. (1999). Non-pathogenic *Escherichia coli* versus mesalazine for the treatment of ulcerative colitis: a randomised trial. *Lancet* 354, 635–639. doi: 10.1016/s0140-6736(98)06343-0
- Roberfroid, M. B. (1993). Dietary fiber, inulin, and oligofructose: a review comparing their physiological effects. *Crit. Rev. Food Sci. Nutr.* 33, 103–104. doi: 10.1080/10408399309527616
- Roy, S., and Dhaneshwar, S. (2023). Role of prebiotics, probiotics, and synbiotics in management of inflammatory bowel disease: Current perspectives. *World J. Gastroenterol.* 29, 2078–2100. doi: 10.3748/wjg.v29.i14.2078
- Sabater-Molina, M., Larqué, E., Torrella, F., and Zamora, S. (2009). Dietary fructooligosaccharides and potential benefits on health. *J. Physiol. Biochem.* 65, 315–328. doi: 10.1007/BF03180584
- Safavi, M., Farajian, S., Kelishadi, R., Mirlohi, M., and Hashempour, M. (2013). The effects of synbiotic supplementation on some cardio-metabolic risk factors in overweight and obese children: a randomized triple-masked controlled trial. *Int. J. Food Sci. Nutr.* 64, 687–693. doi: 10.3109/09637486.2013.775224
- Sanchez, M., Darimont, C., Drapeau, V., Emady-Azar, S., Lepage, M., Rezzonico, E., et al. (2014). Effect of *Lactobacillus rhamnosus* CGMCC1.3724 supplementation on weight loss and maintenance in obese men and women. *Br. J. Nutr.* 111, 1507–1519. doi: 10.1017/s0007114513003875
- Schiffirin, E., Rochat, F., Link-Amster, H., Aeschlimann, J., and Donnet-Hughes, A. (1995). Immunomodulation of human blood cells following the ingestion of Lactic acid bacteria. *J. Dairy Sci.* 78, 491–497. doi: 10.3168/jds.s0022-0302(95)76659-0
- Seo, G. S., and Chae, S. C. (2014). Biological therapy for ulcerative colitis: an update. *World J. Gastroenterol.* 20, 13234–13238. doi: 10.3748/wjg.v20.i37.13234
- Seong, G., Lee, S., Min, Y. W., Jang, Y. S., Park, S. Y., Kim, C. H., et al. (2020). Effect of a synbiotic containing *Lactobacillus paracasei* and *Opuntia humifusa* on a murine model of irritable bowel syndrome. *Nutrients* 12, 3205. doi: 10.3390/nu12103205
- Shahani, K. M., and Chandan, R. C. (1979). Nutritional and healthful aspects of cultured and culture-containing dairy foods. *J. Dairy Sci.* 62, 1685–1694. doi: 10.3168/jds.s0022-0302(79)83481-5
- Sharma, P., Kaur, S., Chadha, B. S., Kaur, R., Kaur, M., and Kaur, S. (2021). Anticancer and antimicrobial potential of enterocin 12a from *Enterococcus faecium*. *BMC Microbiol.* 21, 39. doi: 10.1186/s12866-021-02086-5
- Sharma, M., and Shukla, G. (2016). Metabiotics: One Step ahead of probiotics; an insight into mechanisms involved in anticancerous effect in colorectal cancer. *Front. Microbiol.* 7. doi: 10.3389/fmicb.2016.01940
- Sheoran, P., and Tiwari, S. K. (2019). Enterocin LD3 from *Enterococcus hirae* LD3 causing efflux of intracellular ions and UV absorbing materials in gram negative bacteria. *J. Appl. Microbiol.* 126, 1059–1069. doi: 10.1111/jam.14203
- Sheoran, P., and Tiwari, S. K. (2020). Synergistically-acting Enterocin LD3 and Plantaricin LD4 against gram-positive and gram-negative pathogenic bacteria. *Probiotics Antimicrobial Proteins* 13, 542–554. doi: 10.1007/s12602-020-09708-w
- Shin, H. S., Lee, J. H., Pestka, J., and Ustunol, Z. (2000). Growth and viability of commercial *Bifidobacterium* spp in skim milk containing oligosaccharides and inulin. *J. Food Sci.* 65, 884–887. doi: 10.1111/j.1365-2621.2000.tb13605.x
- Sitonen, S., Vapaatalo, H., Salminen, S., Gordin, A., Saxelin, M., Wikberg, R., et al. (1990). Effect of *Lactobacillus* GG yoghurt in prevention of antibiotic associated diarrhoea. *Ann. Med.* 22, 57–59. doi: 10.3109/07853899009147243
- Silva, N. O. E., Brito, B. B. D., Silva, F. A. F. D., Santos, M. L. C., and Melo, F. F. D. (2020). Probiotics in inflammatory bowel disease: Does it work? *World J. Meta-Analysis* 8, 54–66. doi: 10.13105/wjma.v8.i2.54
- Silveira, D. S. C., Veronez, L. C., Lopes-Junior, L. C., Anatriello, E., Brunaldi, M. O., and Pereira-da-Silva, G. (2020). *Lactobacillus bulgaricus* inhibits colitis-associated cancer via a negative regulation of intestinal inflammation in azoxymethane/dextran sodium sulfate model. *World J. Gastroenterol.* 26, 6782–6794. doi: 10.3748/wjg.v26.i43.6782
- Singh, A., Mahajan, R., Kedia, S., Dutta, A. K., Anand, A., Bernstein, C. N., et al. (2022). Use of thiopurines in inflammatory bowel disease: an update. *Intestinal Res.* 20, 11–30. doi: 10.5217/ir.2020.00155
- Smid, A., Strnisa, L., Bajc, K., Vujic-Podlipic, D., Bogovic Matijasic, B., and Rogelj, I. (2016). Randomized clinical trial: The effect of fermented milk with the probiotic cultures *Lactobacillus acidophilus* La-5[®] and *Bifidobacterium* BB-12[®] and Beneo dietary fibres on health-related quality of life and the symptoms of irritable bowel syndrome in adults. *J. Funct. Foods* 24, 549–557. doi: 10.1016/j.jff.2016.04.031
- Somashekaraiah, R., Mottawea, W., Gunduraj, A., Joshi, U., Hammami, R., and Sreenivasa, M. Y. (2021). Probiotic and antifungal attributes of *Levilactobacillus brevis* MYSN105, isolated from an Indian traditional fermented food pozha. *Front. Microbiol.* 12. doi: 10.3389/fmicb.2021.696267
- Son, S., Koh, J., Park, M., Ryu, S., Lee, W., Yun, B., et al. (2019). Effect of the *Lactobacillus rhamnosus* strain GG and tagatose as a synbiotic combination in a dextran sulfate sodium-induced colitis murine model. *J. Dairy Sci.* 102, 2844–2853. doi: 10.3168/jds.2018-15013
- Sood, A., Mahajan, R., Singh, A., Midha, V., Mehta, V., Narang, V., et al. (2019). Role of faecal microbiota transplantation for maintenance of remission in patients with ulcerative colitis: A pilot study. *J. Crohn's Colitis* 13, 1311–1317. doi: 10.1093/ecco-jcc/jjz060
- Sood, A., Midha, V., Makharia, G. K., Ahuja, V., Singal, D., Goswami, P., et al. (2009). The probiotic preparation, VSL3 induces remission in patients with mild-to-moderately active ulcerative colitis. *Clin. Gastroenterol. Hepatol.* 7, 1202–1209.e1. doi: 10.1016/j.cgh.2009.07.016
- Steinhart, A., Feagan, B. G., Wong, C. J., Vandervoort, M., Mikolainis, S., Croitoru, K., et al. (2002). Combined budesonide and antibiotic therapy for active Crohn's disease: A randomized controlled trial. *Gastroenterology* 123, 33–40. doi: 10.1053/gast.2002.34225
- Su, H., Kang, Q., Wang, H., Yin, H., Duan, L., Liu, Y., et al. (2018). Effects of glucocorticoids combined with probiotics in treating crohn's disease on inflammatory factors and intestinal microflora. *Exp. Ther. Med.* 16, 2999–3003. doi: 10.3892/etm.2018.6559
- Sutas, Y., Hurme, M., and Isolauri, E. (1996). Down regulation of anti-CD3 antibody induced IL-4 production by bovine caseins hydrolysed with *Lactobacillus* GG derived enzymes. *Scandinavian J. Immunol.* 43, 687–689. doi: 10.1046/j.1365-3083.1996.d01-258.x
- Swanson, K. S., Gibson, G. R., Hutkins, R., Reimer, R. A., Reid, G., Verbeke, K., et al. (2020). The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of synbiotics. *Nat. Rev. Gastroenterol. Hepatol.* 17, 687–701. doi: 10.1038/s41575-020-0344-2
- Tajadadi-Ebrahimi, M., Bahmani, F., Shakeri, H., Hadaegh, H., Hijjajafari, M., Abedi, F., et al. (2014). Effects of daily consumption of synbiotic bread on insulin metabolism and serum high-sensitivity C-reactive protein among diabetic patients: A double-blind, randomized, controlled clinical trial. *Ann. Nutr. Metab.* 65, 34–41. doi: 10.1159/000365153
- Tamaki, H., Nakase, H., Inoue, S., Kawanami, C., Itani, T., Ohana, M., et al. (2015). Efficacy of probiotic treatment with *Bifidobacterium longum* 536 for induction of remission in active ulcerative colitis: A randomized, double blinded, placebo controlled multicenter trial. *Digestive Endoscopy* 28, 67–74. doi: 10.1111/den.12553
- Tharmaraj, N., and Shah, N. P. (2009). Antimicrobial effects of probiotic bacteria against selected species of yeasts and moulds in cheese-based dips. *Int. J. Food Sci. Technol.* 44, 1916–1926. doi: 10.1111/j.1365-2621.2009.01986.x
- Tojo, M., Oikawa, T., Morikawa, Y., Yamashita, N., Iwata, S., Satoh, Y., et al. (1987). The effects of *Bifidobacterium breve* administration on *Campylobacter enteritis*. *Pediatr. Int.* 29, 160–167. doi: 10.1111/j.1442-200x.1987.tb00024.x
- Tolonen, A. C., Beauchemin, N., Bayne, C., Li, L., Tan, J., Lee, J., et al. (2022). Synthetic glycans control gut microbiome structure and mitigate colitis in mice. *Nat. Commun.* 13, 1244. doi: 10.1038/s41467-022-28856-x
- Turner, D., Ricciuto, A., Lewis, A., D'Amico, F., Dhaliwal, J., Griffiths, A. M., et al. (2021). STRIDE-II: An update on the selecting therapeutic targets in inflammatory bowel disease (STRIDE) Initiative of the International Organization for the Study of IBD (IOIBD): Determining therapeutic goals for treat-to-target strategies in IBD. *Gastroenterology* 160, 1570–1583. doi: 10.1053/j.gastro.2020.12.031

- Tursi, A., Brandimarte, G., Papa, A., Giglio, A., Elisei, W., Giorgetti, G. M., et al. (2010). Treatment of relapsing mild-to-moderate ulcerative colitis with the probiotic VSL3 as adjunctive to a standard pharmaceutical treatment: A double-blind, randomized, placebo-controlled Study. *Am. J. Gastroenterol.* 105, 2218–2227. doi: 10.1038/ajg.2010.218
- Tzounis, X., Rodriguez-Mateos, A., Vulevic, J., Gibson, G. R., Kwik-Urbe, C., and Spencer, J. P. (2011). Prebiotic evaluation of cocoa-derived flavanols in healthy humans by using a randomized, controlled, double-blind, crossover intervention study. *Am. J. Clin. Nutr.* 93, 62–72. doi: 10.3945/ajcn.110.000075
- Ukwah, B. N., Ezeonu, I. M., and Uguru, J. (2014). Assessment of prebiotic roles of *Musa paradisiaca* (Plantain) peels for controlling spread of *Salmonella* and other infectious disease agents by broilers. *J. Biol. Chem. Res.* 31, 901–911.
- Ungaro, R., Mehandru, S., Allen, P. B., Peyrin-Biroulet, L., and Colombel, J. F. (2017). Ulcerative colitis. *Lancet* 389, 1756–1770. doi: 10.1016/s0140-6736(16)32126-2
- Valcheva, R., Koleva, P., Martinez, I., Walter, J., Ganzle, M. G., and Dieleman, L. A. (2018). Inulin-type fructans improve active ulcerative colitis associated with microbiota changes and increased short-chain fatty acids levels. *Gut Microbes* 10, 334–357. doi: 10.1080/19490976.2018.1526583
- Valcheva, R., Armstrong, H., Kovic, O., Bording-Jorgensen, M., Veniamin, S., Pérez-Muñoz, M. E., et al. (2022). Double blind placebo-controlled trial for the prevention of ulcerative colitis relapses by β -fructan prebiotics: efficacy and metabolomic analysis. Preprint. Available from: medRxiv. doi: 10.1101/2022.01.16.22269376
- Van der Beek, C. M., Dejong, C. H. C., Troost, F. J., Masclee, A. A. M., and Lenaerts, K. (2017). Role of short-chain fatty acids in colonic inflammation, carcinogenesis, and mucosal protection and healing. *Nutr. Rev.* 75, 286–305. doi: 10.1093/nutrit/nuw067
- Van der Meer, R., and Bovee-Oudenhoven, I. (1998). Dietary modulation of intestinal bacterial infections. *Int. Dairy J.* 8, 481–486. doi: 10.1016/s0958-6946(98)00072-7
- Van Laere, K. M. J., Hartemink, R., Bosveld, M., Schols, H. A., and Voragen, A. G. J. (2000). Fermentation of plant cell wall derived polysaccharides and their corresponding oligosaccharides by intestinal bacteria. *J. Agric. Food Chem.* 48, 1644–1652. doi: 10.1021/jf990519i
- Venturi, A., Gionchetti, P., Rizzello, F., Johansson, R., Zucconi, E., Brigidi, P., et al. (1999). Impact on the composition of the faecal flora by a new probiotic preparation: preliminary data on maintenance treatment of patients with ulcerative colitis. *Alimentary Pharmacol. Ther.* 13, 1103–1108. doi: 10.1046/j.1365-2036.1999.00560.x
- Vernero, M., De Blasio, F., Ribaldone, D. G., Bugianesi, E., Pellicano, R., Saracco, G. M., et al. (2020). The usefulness of microencapsulated sodium butyrate add-on therapy in maintaining remission in patients with ulcerative colitis: A prospective observational study. *J. Clin. Med.* 9, 3941. doi: 10.3390/jcm9123941
- Vyas, U., and Ranganathan, N. (2012). Probiotics, prebiotics, and synbiotics: Gut and beyond. *Gastroenterol. Res. Pract.* 2012, 1–16. doi: 10.1155/2012/872716
- Waljee, A. K., Wiitala, W. L., Govani, S., Stidham, R., Saini, S., Hou, J., et al. (2016). Corticosteroid use and complications in a US inflammatory bowel disease cohort. *PLoS One* 11, e0158017. doi: 10.1371/journal.pone.0158017
- Wang, W., Chen, L., Zhou, R., Wang, X., Song, L., and Huang, S. (2014). Increased proportions of *Bifidobacterium* and the *Lactobacillus* group and loss of butyrate-producing bacteria in inflammatory bowel disease. *J. Clin. Microbiol.* 52, 398–406. doi: 10.1128/jcm.01500-13
- Wang, Y. N., Meng, X. C., Dong, Y. F., Zhao, X. H., Qian, J. M., Wang, H. Y., et al. (2019). Effects of probiotics and prebiotics on intestinal microbiota in mice with acute colitis based on 16S rRNA gene sequencing. *Chin. Med. J.* 132, 1833–1842. doi: 10.1097/cm9.0000000000000308
- Wild, G. E., Waschke, K. A., Bitton, A., and Thomson, A. B. (2003). The mechanisms of prednisone inhibition of inflammation in crohn's disease involve changes in intestinal permeability, mucosal TNF α production and nuclear factor kappa B expression. *Aliment Pharmacol. Ther.* 18, 309–317. doi: 10.1046/j.1365-2036.2003.01611.x
- Wildt, S., Nordgaard, I., Hansen, U., Brockmann, E., and Rumessen, J. J. (2011). A randomized double-blind placebo-controlled trial with *Lactobacillus acidophilus* La-5 and *Bifidobacterium animalis* subsp. lactis BB-12 for maintenance of remission in ulcerative colitis. *J. Crohn's Colitis* 5, 115–121. doi: 10.1016/j.crohns.2010.11.004
- Wirngo, F. E., Lambert, M. N., and Jeppesen, P. B. (2016). The physiological effects of dandelion (*Taraxacum officinale*) in type 2 diabetes. *Rev. Diabetic studies: RDS.* 13, 113–131. doi: 10.1900/RDS.2016.13.113
- Wong, W.-Y., Chan, B. D., Leung, T.-W., Chen, M., and Tai, W. C.-S. (2022). Beneficial and anti-inflammatory effects of formulated prebiotics, probiotics, and synbiotics in normal and acute colitis mice. *J. Funct. Foods* 88, 104871. doi: 10.1016/j.jff.2021.104871
- Wong, V. W. S., Wong, G. L. H., Chim, A. M. L., Chu, W. C. W., Yeung, D. K. W., Li, K. C. T., et al. (2013). Treatment of nonalcoholic steatohepatitis with probiotics. A proof-of-concept study. *Ann. Hepatol.* 12, 256–262. doi: 10.1016/s1665-2681(19)31364-x
- Xue, Z., Li, R., Liu, J., Zhou, J., Zhang, X., Zhang, T., et al. (2023). Preventive and synbiotic effects of the soluble dietary fiber obtained from *Lentinula edodes* byproducts and *Lactobacillus plantarum* LP90 against dextran sulfate sodium induced colitis in mice. *J. Sci. Food Agric.* 103, 616–626. doi: 10.1002/jsfa.12173
- Yang, S. C., Chen, J. Y., Shang, H. F., Cheng, T. Y., Tsou, S. C., and Chen, J. R. (2005). Effect of synbiotics on intestinal microflora and digestive enzyme activities in rats. *World J. Gastroenterol.* 11, 7413. doi: 10.3748/wjg.v11.i47.7413
- Yilmaz, I., Dolar, M. E., and Ozpinar, H. (2019). Effect of administering kefir on the changes in fecal microbiota and symptoms of inflammatory bowel disease: A randomized controlled trial. *Turkish J. Gastroenterol.* 30, 242–253. doi: 10.5152/tjg.2018.18227
- Yoo, J. W., Shin, Y. J., Ma, X., Son, Y. H., Jang, H. M., Lee, C. K., et al. (2022). The alleviation of gut microbiota-induced depression and colitis in mice by anti-inflammatory probiotics NK151, NK173, and NK175. *Nutrients* 14, 2080. doi: 10.3390/nu14102080
- Yoshimatsu, Y., Yamada, A., Furukawa, R., Sono, K., Osamura, A., Nakamura, K., et al. (2015). Effectiveness of probiotic therapy for the prevention of relapse in patients with inactive ulcerative colitis. *World J. Gastroenterol.* 21, 5985–5994. doi: 10.3748/wjg.v21.i19.5985
- Zeng, H., Xue, Y., Peng, T., and Shao, W. (2007). Properties of xylanolytic enzyme system in *bifidobacteria* and their effects on the utilization of xylooligosaccharides. *Food Chem.* 101, 1172–1177. doi: 10.1016/j.foodchem.2006.03.019
- Zocco, M. A., Dal Verme, L. Z., Cremonini, F., Piscaglia, A. C., Nista, E. C., Candelli, M., et al. (2006). Efficacy of *Lactobacillus* GG in maintaining remission of ulcerative colitis. *Alimentary Pharmacol. Ther.* 23, 1567–1574. doi: 10.1111/j.1365-2036.2006.02927.x

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
1	DR. SAYABU LAXMAN GAIKWAD	Professor / Principal / Superintendent	RESEARCH	ROLE OF NARIKEL LAVANA IN AMLAPITTA (HYPER ACIDITY)	International	01:00	ROLE OF NARIKEL LAVANA IN AMLAPITTA (HYPER ACIDITY)	9	1	23 - 24	ROLE OF NARIKEL LAVANA IN AMLAPITTA (HYPER ACIDITY)	2020	2278-7348
2			RESEARCH	A CLINICAL MANAGEMENT OF SHUKRAKSHAYA WITH APATYAKARAGHRITA W.S.R. TO OLIGOSPERMIA	International	01:00	INTERNATIONAL JOURNAL OF RESEARCH IN INDIAN MEDICINE	4	3	01-Feb	INTERNATIONAL JOURNAL OF RESEARCH IN INDIAN MEDICINE	2020	2456/4435
3			RESEARCH	TEACHING AND LEARNING METHODS IN AYURVEDA AND ITS CURRENT PERSPECTIVES.	National	01:00	AYURLOG - NATIONAL JOURNAL OF RESEARCH IN AYURVED SCIENCE	8	3	01-Feb	AYURLOG - NATIONAL JOURNAL OF RESEARCH IN AYURVED SCIENCE	2020	2320/7329
4			RESEARCH	AN AYURVED MANAGEMENT OF TRIGEMINAL NEURALGIA	International	01:00	WORLD JOURNAL OF PHARMACEUTICAL RESEARCH	9	7	05-Jun	WORLD JOURNAL OF PHARMACEUTICAL RESEARCH	2020	2277-7105
5			RESEARCH	CONCEPT OF OJAS AND ITS APPLICATION IN RAJAYAKSHMA	International	01:00	WORLD JOURNAL OF PHARMACEUTICAL RESEARCH	9	7	08-Sep	WORLD JOURNAL OF PHARMACEUTICAL RESEARCH	2020	2277-7105
6			RESEARCH	A Clinical Management of Shukrashaya with apatyakaraghrita w.s.r.to oligospermia	International	1 : , 2 : , 3 :	Ayurline: International journal of research in indian medicine	4th	3rd	01-May	Ayurline	2020	2456-4435

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
7			RESEARCH	An open clinical study of jaloukava charana with Abhyantara prayoga of Abhyayadi (haritaki) Ghruta in Sirajaal with Special reference to Episcleritis	International	1 : , 2 :	Vidyasearch	8th	2nd	35 - 40	vidyasearch	2019	2278-7348
8			RESEARCH	EFFICACY OF ANTENATAL MANAGEMENT IN AYURVED DURING 8th AND 9th MONTH OF PREGNANCY	International	1 : , 2 : , 3 :	EFFICACY OF ANTENATAL MANAGEMENT IN AYURVED DURING 8th AND 9th	8	2	3194 - 3195	AYUSHDHARA	2021	2393-9583 (P) 2393-9591
1	DR. RAMAKANT PRADYUMNA AYACHIT	Assistant Professor	RESEARCH	sanskrite vidnyanam	International	01:00	Worldwide International Interdisciplinary Research Journal	1	XXIII	627 - 628	Siddhi Publication	2020	ISSN-2454-7905
1	DR. DIPALI UTTAMRAO SUKE	Assistant Professor	RESEARCH	PROSPECTIVE STUDY OF KAMALA W.S.SR. TO BRIHATRAYEE	International	:	IJAAM	3	6	295 - 301	IJAAM	2015	p-ISSN-2395-3985
2			RESEARCH	CONCEPT OF AMLAPITTA W.S.R. TO MAHASROTAS	International	:	IJAAM	4	6	263 - 268	IJAAM	2016	p-ISSN-2395-3985
3			RESEARCH	THE STUDY OF GRIDHRASI ACCORDING TO CHAKRAPANIDATTA AND ITS MANAGEMENT BY PANCHMULI KASHAY WITH ERAND TAIL	International	:	IJAAM	5	4	178 - 183	IJAAM	2017	p-ISSN-2395-3985

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
4			RESEARCH	AHAR VIDHI: CONCEPT OF FOOD INTAKE IN AYURVEDA WITH COMPARISON TO PRESENT ERA	International	:	INTERNATIONAL RESEARCH JOURNAL OF INDIA	1	7	25 - 30	IRJI	2016	ISSN NO-2454-8707
5			RESEARCH	CONCEPT OF VIRUDDHA AHAR W.S.R. TO ABHISHYANDI GUNA	International	:	WORLD JOURNAL OF PHARMACEUTICAL RESEARCH	6	3	1678 - 1685	WJPR	2017	ISSN 2277-7105
6			RESEARCH	THE MAHASNEHA MATRABASTI IN ASTHI-MAJJA AVRIT VATA W.S.R. TO COMPRESSIVE NEUROPATHY - A CASE STUDY	International	1 : DIPALI UTTAMRAO SUKE, 2 :	INTERNATIONAL JOURNAL OF AYURVEDA & ALTERNATIVE MEDICINE	8	1	31 - 34	IJAAM	2020	p-ISSN - 2395 - 3985, e-ISSN - 2348 -0173
			RESEARCH	THE STUDY OF AHAR MATRA W.S.R. TO JATHARAGNI AND ITS IMPORTANCE IN CASE HISTORY	International	1 : Pathak Radhika Govind, 2 : Suke	IAMJ	p5	6	3109 - 3112	IAMJ	2021	ISSN: 2320-5091
7			RESEARCH	Comparative study of santarpanjanya vyadhi hetu w.s.r to Bruhatrayee	International	1 : Dr Dipali Uttamrao Suke	IJRAR	10	3	578 - 583	IJPUBLICATION	2023	E-ISSN:2348-1269,P-ISSN:2349-5138
1	DR. HARSHA SHANKAR GAMBHIRE	Associate Professor	RESEARCH	Study of Eye Anatomy (Netra Sharir) in Computer Vision Syndrome	International	:	International Journal of Advanced Research (IJAR)	7	3	151 - 153	International Journal of Advanced Research (IJAR)	2019	2320-5407

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
2			RESEARCH	Study of correlation between Ayurvedic concept of kshetra as garbhsambhav samugri & modern concept of endometrium.	International	:	International Journal of Scientific Research	8	8	46 - 48	International Journal of Scientific Research	2019	2277-8179
3			RESEARCH	Role of Ayurveda in obesity	National	:	Sankalpa Nimacon - 2019	1	38	14 - 15	Wizcraft Publications & Distribution	2019	978-81-940962-0-7
4			RESEARCH	EFFECTS OF MODERN LIFESTYLE ON KSHETRA : A LITERATURE REVIEW	International	:	INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH (IJSR)	8	10	55 - 56	INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH (IJSR)	2019	2277 - 8179
5			RESEARCH	Effects of changing Lifestyle on Annavaahasrotas: A Literature Review	International	:	International Journal of Trend in Scientific Research and Development (IJTSRD)	4	1	125 - 127	IJTSRD Publication	2019	2456-6470
6			RESEARCH	MEDOVAHA SROTAS & PANCHKARMA CHIKITSA	International	01:00	International Journal of Scientific Research (IJSR)	8	11	22 - 23	International Journal of Scientific Research (IJSR)	2019	2277-8179
7			RESEARCH	Treating Artavkshaya by Lifestyle Changes: A Case Study	International	:	International Journal of Multidisciplinary Educational Research (IJMER)	8	12(2)	06-Oct	Sucharitha Publications	2019	2277-7881

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
8			RESEARCH	Anatomical Abnormalities of Kshetra (Uterus): A Literature Review	International	:	International Journal of Multidisciplinary Educational Research (IJMER)	9	1(7)	73 - 79	Sucharitha Publications	2020	2277-7881
9			RESEARCH	An observational study of endometrial pattern in implantation	International	:	INTERNATIONAL JOURNAL OF AYURVEDIC MEDICINE	13	4 (2022): October - December 2022	921 - 924	AYURVEDA SAHITI PRABHA , 1-2-288-23, SBH COLONY, HYDERABAD, ANDHRA PRADESH, INDIA, 500	2023	0976-5921
10			RESEARCH	Treating thin endometrium by Ayurvedic Medications: A Case Study	International	:	INTERNATIONAL JOURNAL OF AYURVEDIC MEDICINE	13	4 (2022): October - December 2022	1102 - 1107	AYURVEDA SAHITI PRABHA , 1-2-288-23, SBH COLONY, HYDERABAD, ANDHRA PRADESH, INDIA, 500	2023	0976-5921
11			RESEARCH	Treating Bilateral Cornual Tubal Block by Ayurvedic Medications: A Case Study	International	:	INTERNATIONAL JOURNAL OF LIFE SCIENCE AND PHARMA RESEARCH	13	1, January 2023	245 - 252	INT JOURNAL LIFESCIENCE & PHARMA RESEARCH, PLOT NO 10, 2 MAIN RD, RENGANAGAR NEAR TO OLD	2023	2250-0480
12			RESEARCH	Evaluation of Heel pain in association with calcaneal spur length and angle	National	:	The journal of oriental Research Madras (Print only)	XCIII-XV	Nov-22	01-Nov	The Kuppuswami Sastri Research Institute, Madras	2022	0022-3301
13			RESEARCH	Treating infertility due to thin endometrium by ayurvedic medications: A case study	National	:	The journal of oriental Research Madras (Print only)	XCIII-XV	Nov-22	Dec-22	The Kuppuswami Sastri Research Institute, Madras	2022	0022-3301

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)	
1	DR. SUHAS DNYANOBA MOTE	Assistant Professor	RESEARCH	A BRIEF REVIEW ON ORGANOPHOSPHORUS POISONING IN AYURVEDA	International	1 : , 2 :	Unique Journal of Medical and Dental Sciences	9	1	05-Sep	Unique Journal of Medical and Dental Sciences	2021	2347-5579	
1	DR. PRIYADARSHANI PANDURANG RAJAGE	Associate Professor	RESEARCH	Role of ayurveda in neurological disorder	National	1 : Dr. Priyadarshani Pandurang	Sankalpa Nimacon 2019	1	1	49 - 50	Wizcraft Publications and Distributions	2019	ISBN 978-81-940962-0-7	
2			RESEARCH	ROLE OF AYURVEDIC TRADITIONAL CONCEPTS FOR PREVENTION OF LIFESTYLE DISORDERS	International	1 : Dr. RAJAGE PRIYADARSHANI PANDUR	CONTEMPORARY RESEARCH IN INDIA	-	-	39 - 41	Wizcraft publication	2020	ISSN 2231-2137	
3			RESEARCH	Role of prakruti in clinical diagnosis	International	1 : Dr. Rajage Priyadarshani Pandura	Journal of emerging technologies and innovative research JETIR	August 2021 volume 8	8	0 - 0	Jetir	2021	ISSN 2349-5162	
4			RESEARCH	Ayurvedic review on concept of agni	International	1 : Dr. Rajage Priyadarshani Pandura	International journal of All Research Education and Scientific Methods	08-Aug-21	9	08-Aug-21	2264 - 2267	IJARESM	2021	ISSN 2455-6211
5			RESEARCH	Application of Sarata Parikshan	International	1 : Dr. Rajage Priyadarshani Pandura	Journal of Emerging Technologies and innovative research JETIR	08-Aug-21	8	8	0 - 0	JETIR publication	2021	ISSN 2349 5162

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
1	DR. RAVIKIRAN ACHYUT DHENGALE	Assistant Professor	RESEARCH	Comparative Study Between Nidhrabhrunsha & Dwidoshaja Prakruti W.s.r. in-somania	International	1 : , 2 :	IAMJ	7	12	3207 - 3212	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	2020	2320/5091
2			RESEARCH	STUDY THE EFFICACY OF CHATURJATADI CHURN IN THE MANAGEMENT OF BALKRUSHA W.S.R. TO UNDERVEIGHT CHILDREN	International	1 : , 2 : , 3 :	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	7	12	2170 - 2177	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	2019	2320/5091
1	DR. SULAKSHANA BABANRAO VARPE	Associate Professor	RESEARCH	ROLE OF TAKRARISTA IN GRAHANI	International	1 : , 2 :	DEERGHAYU INTERNATIONAL	33	129	61 - 67	DEERGHAYU PUBLICATION	2017	0970-3381
2			RESEARCH	COSMETIC APPROACH TO HAIR IN AYURVEDA	International	1 : , 2 :	DEERGHAYU INTERNATIONAL	33	130	115 - 121	DEERGHAYU INTERNATIONAL	2017	0970-3381
3			RESEARCH	STUDY OF USES OF CHANDRAPRABHA VATI IN GYNACOLOGICAL DISORDERS	International	1 : , 2 : , 3 :	DEERGHAYU INTERNATIONAL	33	130	145 - 153	DEERGHAYU INTERNATIONAL	2017	0970=3381
4			RESEARCH	Pharmaceutical and analytical study of dhatri avleha	International	1 : , 2 : , 3 :	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	5	10	3824 - 3833	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	2017	2320 5091

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
5			RESEARCH	PHARMACEUTICAL AND ANALYTICAL STUDY OF DHATRI AVLEHA	International	1 : DR SULAKS HANA VARPE, 2 : DR	INTERNATIONAL AYURVEDIC MEDICAL JOOURNAL	5	10	3824 - 3833	IAMJ	2017	2320 5091
1	DR. MEERA DATTATRAY SURYAWANSHI	Assistant Professor	RESEARCH	HAIRFALL (KHALITYA) A CRITICAL REVIEW, MANAGEMENT & FUTURE ASCEPTS	National	1 : , 2 : , 3 :	AYURVED AN ETERNAL SCIENCE	8	11	38 - 41	AYURVED AN ETERNAL SCIENCE	2020	2231/2137
1	DR. SAMPADA CHETAN SAHASTRABUDDHE	Associate Professor	RESEARCH	AYURVEDIC MANAGEMENT IN OSTEOMYELITIS : A CASE STUDY.	International	1 : , 2 :	International Journal of Ayurveda and Pharma Research (IJAPR)	8	6	71 - 74	Mahadev Publications, R.R. Dist. Telangana, India.	2020	2322 - 0910
2			RESEARCH	WORM INFESTATION [KRUMI] and ITS AYURVEDA ETIOPATHOGENESIS	International	1 : DR SAMPADA A CHETAN SAHASR	INTERNATIONAL JOURNAL OF INDIAN MEDICINE	2	7	27 - 35	AYURVEDA RESEARCH & CAREER ACADEMY [ARCA]	2021	2582:7634
3			RESEARCH	A REVIEW OF SHWASANA-RESPIRATION IN VEDIC TEXTS	National	1 : DR SAMPADA A CHETAN SAHASR	AYURVED PATRIKA	75	7	46 - 49	NATIONAL SCIENTIFIC JOURNAL OF AYURVED	2022	2278-0726
1	DR. PAKANIKAR SATISH LIMBAJI	Associate Professor	RESEARCH	A CRITICAL REVIEW ON MEDOHAR AND LEKHANIYA DRAVYAS	International	:	WORLD JOURNAL OF PHARMACEUTICAL AND MEDICAL RESEARCH	8	92,022	57 - 59	WORLD JOURNAL OF PHARMACEUTICAL AND MEDICAL RESEARCH	2022	2455-3301

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
2			RESEARCH	ROLE OF RASAYAN DRUGS IN AYURVEDA :A REVIEW	International	1 : DR.SATISH LIMBAJI PAKNIKA	INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS (IJCRT)	10	Aug-22	109 - 115	INTERNATIONAL JOURNAL OF CREATIVE RESEARCH THOUGHTS(IJCRT)	2022	2320-2882
1	DR. SHINDE SHREYASH AJIT	Assistant Professor	RESEARCH	A CRITICAL REVIEW ON MEDOHAR AND LEKHANIYA DRAVYAS	International	1 : , 2 :	WORLD JOURNAL OF PHARMACEUTICAL AND MEDICAL RESEARCH	8	9, 2022	57 - 59	WORLD JOURNAL OF PHARMACEUTICAL AND MEDICAL RESEARCH	2022	ISSN 2455-3301
2			RESEARCH	ROLE OF RASAYAN DRUGS IN AYURVEDA: A REVIEW	International	1 : , 2 :	International Journal of Creative Research Thoughts (IJCRT)	10	Aug-22	0 - 0	International Journal of Creative Research Thoughts (IJCRT)	2022	ISSN:2320-2882
1	DR. MANASI MAHESH KULKARNI	Professor	RESEARCH	Role of Shatapushpadi Basti in Abijotsaraga with Respect To Anovulatory Cycle	National	1 : , 2 : , 3 :	Sankalpa Nimacon 2019	10	5	41 - 42	Sankalpa Nimacon 2019	2019	978-81-940962-0-7
2			RESEARCH	ROLE OF AYURVEDA FOR MANAGEMENT OF ANKYLOSING SPONDYLITIS- A CASE STUDY	International	:	INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH	8/Issue-12/December-2019	12	30 - 32	INDIAN SOCIETY FOR HEALTH AND ADVANCED RESEARCH	2019	ISSN NO. - 2277-8179
3			RESEARCH	A CLINICAL APPROACH OF AYURVEDA RELATED TO ATYAMBUPAN WITH RESPECT TO OVERHYDRATION	National	:	INDIAN JOURNAL OF APPLIED RESEARCH	9	12	45 - 46	INDIAN JOURNAL OF APPLIED RESEARCHIS	2019	ISSN NO.- 2249-555X

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
4			RESEARCH	Ayurvedic Management of Mukhapaka w.s.r. to Stomatitis : A Case Study	International	1 : Sulbha kalame, 2 : Manasi	Ayurline: International Journal of Research in Indian Medicine	05th	3rd	0 - 0	Ayurline IJ -RIM	2021	2456-4435
5			RESEARCH	Role of Ayurveda for the management of COVID-19 A Case Study	International	1 : Dr.Mana si Kulkarni, 2 :	International Journal of Current Advanced Research	10	10(B)	25370 - 25373	International Journal of Current Advanced Research	2021	2319-6475
6			RESEARCH	Shardiya Navratra-Aamalki Rasayan Mahotsav	National	1 : Dr.Mana si Kulkarni	National Scientific Journal of Ayurved	75	3	14 - 19	National Scientific Journal of Ayurveda	2021	2278-0726
1	DR. NURJAHAN TAJODDIN MUJAWAR	Assistant Professor	RESEARCH	Role of Ayurveda in Obesity	National	1 : , 2 : , 3 :	Sankalpa Nimacon 2019	1	1	31 - 32	Wizcraft Publications and Distribution	2019	ISBN - 978-81-940962-0-7
2			RESEARCH	Role of Ayurveda for management of covid 19 A case study	International	1 : Dr.Mana si Mahesh Kulkarni, 2 :	International Journal of current advanced research	10	10 (B)	25370 - 25373	International journal of current advanced research	2021	2319-6475
3			RESEARCH	Ayurvedic management of mukhpaka w.s.r. to stomatitis	International	1 : Dr Manasi M. Kulkarni, 2 : Dr.	International Journal of research in indian medicine	05 th	3rd	01-May	International journal of research in Indian medicine	2021	2456-4435

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
1	DR. PRADEEP UTTAM JADHAV	Associate Professor	RESEARCH	In vivo study of efficacy of Arka (Calotropis procera, Ait.) moola swaras as first aid measure in snake venom poisoning.	International	1 : Dr Jadhav Pradeep Uttam, 2 : Dr	International Journal of Multidisciplinary Health Sciences	II	1	01-Sep	Mr Sanjay Jadhav, Prakash Shikshan Mandal, Islampur	2016	2394-9406
2			RESEARCH	ARKA (Calotropis procera): A Review	International	1 : Dr Jadhav Pradeep Uttam, 2 : Dr	International journal of Multidisciplinary Health Sciences	IV	1	01-Nov	Mr Sanjay Jadhav, Prakash Shikshan Mandal, Islampur	2018	2394-9406
3			RESEARCH	ASHWAGANDHA (Withania somnifera): A Review	International	1 : Dr Jadhav Pradeep Uttam, 2 : Dr	International Journal of Multidisciplinary Health Sciences	IV	1	01-Nov	Mr Sanjay Jadhav, Prakash Shikshan Mandal, Islampur	2018	2394-9406
4			RESEARCH	In vivo study of role of Ashwagandha (Withania somnifera) root churna as first aid measure in snake venom poisoning.	International	1 : Dr Jadhav Pradeep Uttam, 2 : Dr V.P.	International Journal of Ayurveda & Alternative Medicine.	7	4	132 - 139	Dr. Swati Bhingare, Bhandup, Mumbai.	2019	2348-0173
5			RESEARCH	ROLE OF AYURVEDA IN OBESITY: A CASE STUDY	International	1 : Dr Pradeep Uttam Jadhav, 2 : Dr	International Journal of Ayurveda & Alternative Medicine	7	4	143 - 147	Dr. Swati Bhingare, Bhandup, Mumbai.	2019	2348-0173
6			RESEARCH	STUDY OF THE EFFICACY OF DHANYAKA (CORIANDRUM SATIVUM) WITH SITA (SUGAR CANDY) AND DADHI (CURD) IN JAYPAL (CROTON TIGLIUM)	International	1 : Dr Pachkawade Shubhangi	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL (IAMJ)	8	5	3453 - 3461	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL (IAMJ)	2020	2320 5091

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
7			RESEARCH	A REVIEW ARTICLE ON JAYPAL-CROTON TIGLIUM	International	1 : Dr Jadhav Pradeep Uttam, 2 : Dr	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL (IAMJ)	VIII	6	3714 - 3720	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL (IAMJ)	2020	2320 5091
8			RESEARCH	AYURVEDIC MANAGEMENT IN OSTEOMYELITIS: A CASE STUDY	International	1 : Dr Sahasra budhe Sampada, 2 : Dr	International Journal of Ayurveda and Pharma Research (IJAPR)	8	6	71 - 74	Mahadev Publications, R.R. Dist, Telangana, India	2020	2322 - 0910
9			RESEARCH	Conceptual study on Vajikaran in Ayurveda	International	1 : Prashant Rajaram Shirke, 2 : Jadhav	Journal of Ayurveda and Integrated Medical Sciences (JAIMS)	7	1	240 - 243	JAIMS	2022	ISSN 2456-3110
10			RESEARCH	Ayurvedic treatment in the management of diabetic foot ulcer- A case study	International	1 : Dr Pradeep Uttam Jadhav	World Journal of Pharmaceutical Research (WJPR)	12	20	757 - 762	World Journal of Pharmaceutical Research (WJPR)	2023	2277-7105
1	DR. SHUBHANGI TUKARAM PACHKAWADE	Assistant Professor	RESEARCH	STUDY OF THE EFFICACY OF DHANYAKA (CORIANDRUM SATIVUM) WITH SITA (SUGAR CANDY) AND DADHI (CURD) IN JAYPAL (CROTON TIGLIUM)	International	:	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	8	5	3454 - 3461	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	2020	2320-5091
2			RESEARCH	A REVIEW ARTICLE ON JAYPAL-CROTON TIGLIUM	International	:	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	8	6	3715 - 3720	IAMJ	2020	2320 5091

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
3			RESEARCH	THE MAHASNEHA MATRABASTI IN ASTHI-MAJJA AVRIT VATA W.S.R. TO COMPRESSIVE NEUROPATHY-A CASE STUDY	International	1 : Dipali U. Suke, 2 : Shubhangi T.	INTERNATIONAL JOURNAL AND ALTERNATIVE MEDICINES	8	1/ JAN-FEB- (2020)	31 - 34	INTERNATIONAL JOURNAL AND ALTERNATIVE MEDICINES	2020	eISSN-2348-0173
1	DR. RAVIKIRAN RAM BHOSALE	Professor	RESEARCH	Multimodal Ayurvedic Management of Kashtartava (Dysmenorrhoea) A Case Series	National	:	Ayurlog National Journal of Research in Ayurved Science	8	6	01-Jul	Ayurlog National Journal of Research in Ayurved Science	2020	2320-7329
2			RESEARCH	A Clinical Study to Assess Efficacy of Bilva Majja Churna With Lajaambu Anupana in Garbhini Chharbi (Emesis Gravidarum)	International	1 : , 2 :	Ayurline International Journal of Research in Indian Medicine	5	1	01-Jul	Ayurline International Journal of Research in Indian Medicine	2021	2456-4435
3			RESEARCH	Significance of Sanskara in Sneha Kalpana A Review	National	1 : , 2 :	Ayurlog National Journal of Research in Ayurved Science	9	1	01-Oct	Ayurlog National Journal of Research in Ayurved Science	2021	2320-7329
4			RESEARCH	A CRITICAL REVIEW OF GARBHINI CHHARDI (Emesis gravidarum)	International	1 : Dr. Ravikiran Ram Bhosale, 2 : Dr. Garthe L.S, 2 : Dr. Ravikira	International Journal of Ayurveda & Alternative Medicine	8	4	163 - 165	IJAAM	2020	2395-3985
5			RESEARCH	Clinical Efficacy of local Guggulu Dhoopana in the management of Episotomy wound	National	1 : Dr. Garthe L.S, 2 : Dr. Ravikira	Ayurlog National Journal of Research in Ayurved Science	9	9	01-Feb	Ayurlog National Journal of Research in Ayurved Science	2021	2320-7329

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
1	DR. SULBHA DNYANOBA KALME	Professor	RESEARCH	An Ayurvedic Classical Approach Towards Pcos and Its Management A Literature Review	National	1 : , 2 : , 3 :	Sankalpa Nimacon - 2019	1	1	27 - 28	Wizcraft Publication and Distribution	2019	ISBN - 978-81-940962-0-7
2			RESEARCH	Ayurvedic Approach on Neutraceuticals w.s.r.to Rasayana Therapy	International	01:00	VIDYASEARCH	8th	4th	48 - 54	vidyasearch	2020	2278-7348
3			RESEARCH	Computer Vision Syndrome in Children & its Management : An Ayurvedic Approach	National	1 : , 2 :	Ayurlog : NJ-RAS	8TH	5TH	01-Apr	AYURLOG	2020	2320-7329
4			RESEARCH	Comparative study of role of Yashtimadhu siddha ghruta Tarpan & Anutail Nasya in Shushkakshipaka	International	1 : Dr.Ashwini S Gaikwad , 2 : Dr. - - - - -	ayurline	5th	1st	01-Nov	Internatinal journal of research in indian medicine	2021	2456-4435
5			RESEARCH	Ayurvedic management of mukhapaka w.s.r. to Stomatitis: A case study	International	1 : , 2 : , 3 :	Ayurline	5th	3rd	01-May	Ayurline international journal of research in indian medicine	2021	2456-4435
6			RESEARCH	Role of Ayurveda for the management of Covid-19 A case study	International	1 : Dr.Mansi kulkari, 2 : Dr.Sulbh	International journal of Current Advanced Research	10	10(B)	25370 - 25373	International Journal of Current Advance Research	2021	2319-6475

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
1	DR. PALLAVI MORESHWAR KANITKAR	Associate Professor	RESEARCH	Neem(Azadiracta indica)a possible silver bullet in Covid 19	International	1 : Pallavi Moreshwar Kanitkar,	International journal of research in pharmaceutical science	11	4	1105 - 1109	IJRPS	2020	0975-7538
			RESEARCH	Neem (Azadiracta indica) a possible Silver bullet in Covid 19	International	1 : Pallavika nitkar, 2 : Neeta Deshpan	International journal of research in pharmaceutical science	11	4	1105 - 1109	IJRPS	2020	BBU3MZ
1	DR. MAHESHKUMAR BHARAT RAUT	Assistant Professor	RESEARCH	Study of Ashmari in Ayurvedic and modern point of view	International	1 : Dr. Mahesh kumar Bharat Raut	International Journal of Creative Research Thoughts (IJCRT)	11	11	01-May	International Journal of Creative Research Thoughts (IJCRT)	2023	2320-2882
1	DR. SANDHYA SURESH MANE	Assistant Professor	RESEARCH	Clinical Assessment of Jalaukavcharan (LeechTherapy) in Management of Sirakoutilyajanya Shotha -A Case Report	International	1 : Dr.Sandhya Suresh Mane.	International Ayurveda Medical Journal	VIII	VI	3820 - 3826	IAMJ	2020	2320-5091 /978-1-635354805
2			RESEARCH	A Case Study-Management of Medovrudhi With Special reference to Hyperlipidemia by Ayurvedic Medicine.	International	1 : Dr.Sandhya Suresh Mane	International journal of Ayurvedic and Herbal Medicine	10	5	3839 - 3843	IJAHM	2020	2249-5746
1	DR. VAISHALI PRASHANT MOHIRE	Associate Professor	RESEARCH	A CASE STUDY OF SHIRODHARA WITH YOGNIDRA IN VATAJ SHIRSHOOL (HEADACHE)	National	01:00	INDIAN JOURNAL OF APPLIED RESEARCH	9	12-Dec-19	70 - 71	INDIAN JOURNAL OF APPLIED RESEARCH	2019	2249- 555X DOI : 10.36106/ijar

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
2			RESEARCH	A CASE STUDY OF SHIRODHARA IN TINNITUS	International	01:00	INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH	8	12-Dec-19	70 - 71	INTERNATIONAL JOURNAL OF SCIENTIFIC RESEARCH	2019	2277-8179 DOI 10.36106/ijsr
3			RESEARCH	EFFECT OF JATIPATRA KAVAL (CHARVAN) IN MUKHAPAKA (STOMSTITIS WITH APHTHOUS ULCER)	International	1 : VD. VAISHALI MOHIRE (INTERNATIONAL JOURNAL OF MULTIDISCIPLINARY EDUCATIONAL RESEARCH	9	9 (7) SEPTEMBER 2020	131 - 134	SUCHARITHA PUBLICATIONS , VISAKHAPATANAM, ANDHRA PRADESH , INDIA	2020	2277-7881
4			RESEARCH	A CASE STUDY OF NASYAKARMA IN NASAPRATINAHA - NASAL OBSTRUCTION	International	01:00	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	VIII	IV	3411 - 3413	INTERNATIONAL AYURVEDIC MEDICAL JOURNAL	2020	2320-5091
5			RESEARCH	LOCAL PROCEDURE IN URDHVAJATRUGAT VIKARA, A REVIEW ARTICLE ON "KARNPOORNA"	International	01:00	INTERNATIONAL JOURNAL OF MULTIDISCIPLINARY EDUCATIONAL RESEARCH	10	1(1)	112 - 113	SUCHARITHA PUBLICATIONS	2021	2277-7881
1	DR. ASHWINI SAYABU GAIKWAD	Assistant Professor	RESEARCH	Coparetive study of roll of shatavari - madhuk siddh ghрут tarpan and abhyantarpan in shushkashipak	International	1 : Dr.Ashwini Sayabu Gaikwad	IJRAMS	1	25	125 - 129	IJRAMS	2018	IJRMAS 181-36
2			RESEARCH	An Open Clinical Study of JalokavaCharana with Abhyantara Prayoga of Abhyayadi (Haritaki) Ghruta In Sirajaal with Special Referemnce to Episcleritis	International	1 : , 2 :	VIDYASEARCH	8TH	2278-7348	35 - 40	VIDYASEARCH	2019	2278-7348

Sr. No.	Teacher Name	Designation	Article / Book Type	Title Of Article / Book	Article Type	Name Of Authors	Name Of Journal	Volume No.	Issue No.	Page No.	Name Of Publication	Year Of Publication	Publication Number (ISSN No. / ISB No.)
3			RESEARCH	Computer Vision Syndrome in Children & its Management	National	1 : , 2 :	National Journal of Research in Ayurved Science	8th	5th	01-Apr	AYURLOG	2020	2320-7329
4			RESEARCH	Comparative study of role of Yashthimadhu Siddha Ghruta Tarpan and Anutail Nasya in Sushkakshipaka	International	1 : , 2 :	AYULINE	5TH	1ST	01-Nov	AYURLINE	2021	2456-4435
5			RESEARCH	Ayurvedic management of mukhapaka w.s.r. to stomatitis : A case study	International	1 : , 2 : , 3 :	Ayurline	5th	3rd	01-May	Ayurline	2021	2456-4435

Sr.No.	Name of Teacher	Name Of Research Article	Year Of Publication
1	Dr. RAJAGE PRIYADARSHANI PANDURANG	ROLE OF AYURVEDIC TRADITIONAL CONCEPTS FOR PREVENTION OF LIFESTYLE DISORDERS	2020
2	DR.SAMPADA CHETAN SAHASTRABODDHE	AYURVEDIC MANAGEMENT IN OSTEOMYELITIS : A CASE STUDY.	2020
3	DR.ASHWINI SAYABU GAIKWAD	Computer Vision Syndrome in Children & its Management	2020
4	DR.DIPALI UTTAMRAO SUKE	THE MAHASNEHA MATRABASTI IN ASTHI-MAJJA AVRIT VATA W.S.R. TO COMPRESSIVE NEUROPATHY - A CASE STUDY	2020
5	DR. HARSHA SHANKAR GAMBHIRE	Anatomical Abnormalities of Kshetra (Uterus): A Literature Review	2020
6	DR. MANASI MAHESH KULKARNI	A CLINICAL APPROACH OF AYURVEDA RELATED TO ATYAMBUPAN WITH RESPECT TO OVERHYDRATION	2019
7	DR. NURJAHAN TAJODDIN MUJAV	Role of Ayurveda in Obesity	2019
8	DR.PALLAVI MORESHWAR KANIT	Neem (Azadiracta indica) a possible Silver bullet in Covid 19	2020
9	DR. PRADEEP UTTAM JADHAV	STUDY OF THE EFFICACY OF DHANYAKA (CORIANDRUM SATIVUM) WITH SITA (SUGAR CANDY) AND DADHI (CURD) IN JAYPAL (CROTON TIGLIUM) POISONING IN ALBINO MICE	2020
10	DR. RAMAKANT PRADYUMNA AYACHIT	sanskrite vidnyanam	2020
11	DR. SANDHYA SURESH MANE	A Case Study-Management of Medovrudhi With Special reference to Hyperlipidemia by Ayurvedic Medicine.	2020
12	DR. SAYABU LAXMAN GAIKWAD	A Clinical Management of Shukrashaya with apatyakaraghrita w.s.r.to oligospermia	2020
13	DR. SHUBHANGI TUKARAM PACHKAWADE	A REVIEW ARTICLE ON JAYPAL-CROTON TIGLIUM	2020
14	DR. SULBHA DNYANOBA KALME	Ayurvedic Approach on Nutraceuticals w.s.r.to Rasayana Therapy	2020
15	DR. VAISHALI PRASHANT MOHIRE	A CASE STUDY OF NASYAKARMA IN NASAPRATINAHA - NASAL OBSTRUCTION	2020